

January 1990

The genera of Arundinoideae (Gramineae) in the southeastern United States

Gordon C. Tucker

Eastern Illinois University, gctucker@eiu.edu

Follow this and additional works at: http://thekeep.eiu.edu/bio_fac



Part of the [Biology Commons](#)

Recommended Citation

Tucker, Gordon C., "The genera of Arundinoideae (Gramineae) in the southeastern United States" (1990). *Faculty Research & Creative Activity*. 180.

http://thekeep.eiu.edu/bio_fac/180

This Article is brought to you for free and open access by the Biological Sciences at The Keep. It has been accepted for inclusion in Faculty Research & Creative Activity by an authorized administrator of The Keep. For more information, please contact tabruns@eiu.edu.

JOURNAL
OF THE
ARNOLD ARBORETUM

VOLUME 71

APRIL 1990

NUMBER 2

THE GENERA OF ARUNDINOIDEAE (GRAMINEAE)
IN THE SOUTHEASTERN UNITED STATES^{1, 2}

GORDON C. TUCKER³

Subfamily ARUNDINOIDEAE Tateoka, Jour. Jap. Bot. 32: 277. 1957, "Arun-
doideae."

Perennial or annual, small to very large herbaceous plants of wetlands, wood-
lands, and lowland and montane grasslands [semideserts]. Rhizomes often
present. Stems erect or spreading (stolons sometimes present); nodes solid,

¹Prepared for the Generic Flora of the Southeastern United States, a long-term project made possible by grants from the National Science Foundation and at this writing supported by BSR-8716834 (Norton G. Miller, principal investigator), under which this account was prepared, and BSR-8717333 (Carroll E. Wood, Jr., principal investigator). This treatment, 132nd in the series, follows the format established in the first paper (Jour. Arnold Arb. 39: 296-346. 1958) and continued to the present. The area covered by the Generic Flora includes North and South Carolina, Georgia, Florida, Tennessee, Alabama, Mississippi, Arkansas, and Louisiana. The descriptions are based primarily on the plants of this area, with information about extraregional members of a family or genus in brackets. Those references I did not verify are marked with asterisks.

I have continued to enjoy working with Norton Miller and Carroll Wood on the Generic Flora project and I thank them for their interest and advice. William J. Crins, Ihsan A. Al-Shehbaz, and Barbara Nimblett have helped frequently. Mary E. Barkworth (Utah State University) and Stephen J. Darbyshire (Agriculture Canada, Ottawa) reviewed the manuscript and made numerous valuable suggestions. Elizabeth B. Schmidt and Stephen A. Spongberg improved the final manuscript with their editorial skills. Thanks are extended to the staffs of the New York State Library (especially Alta Beach, Senior Librarian) and the Botany Libraries of Harvard University (especially Geraldine C. Kaye, Librarian) for providing many references. I thank the curators of the following herbaria who have sent specimens or provided access to collections and hospitality during my visits: A, ALU, CCNL, CONN, DUKE, ECON, GH, NEBC, NYS, TRT, UNA, US.

Contribution number 625 of the New York State Science Service.

²The illustrations (by Karen Stoutsenberger from dissections by Carroll Wood) are rearrangements of parts of FIGURES 3, 5, and 7 in C. S. Campbell's account of the family Gramineae, its subfamilies, and tribes in the southeastern United States (Jour. Arnold Arb. 66: 123-199. 1985).

³Biological Survey, New York State Museum, Albany, New York 12230.

glabrous, pubescent, or puberulent. Leaves several, basal and cauline, or cauline only; sheath margin open; blade linear to narrowly lanceolate; bundle sheaths double; chlorenchyma not radiate (except in *Aristida* and *Neyraudia*); silica bodies of various shapes, longitudinally oriented; stomata not dominating intercostal zones, subsidiary cells domed. Inflorescences terminal or terminal and axillary (axillary cleistogamous ones wholly enclosed within sheaths in *Danthonia* and *Nassella*). Spikelets few to very numerous, laterally flattened to subcylindrical, 1- to several-flowered; articulation above the glumes. Glumes 2, subequal to decidedly unequal, lanceolate, equaling or shorter than the lemmas; lemmas conduplicate, involute, or convolute, membranaceous (or indurate). Paleas shorter and narrower than the lemmas. Lodicules [3] 2 (or lacking), oblong or obtruncate, entire or shallowly emarginate, the margin sometimes ciliate. Stamens 1 or 3; filaments slender; anthers ellipsoid to linear. Ovaries ellipsoid to cylindrical. Fruit a caryopsis [achene], cylindrical or flattened, more or less clasped by persistent lemma and palea. Pericarp adnate to [free from] the seed. Hilum ovate-oblong to linear. Base chromosome numbers 9, 10, 11, 12. (Including Aristideae C. E. Hubb., Centothecoideae Soderstrom, Danthoniaceae Zotov, Stipeae Dumort.) TYPE GENUS: *Arundo* L.

A subfamily of about 60 genera and some 1400 species, in five tribes, distributed worldwide but with the greatest diversity, both in species and genera, in the Temperate Zone of the Southern Hemisphere. The Arundinoideae are represented in the Southeast by seven genera in four tribes (Aristideae C. E. Hubb., Arundineae Dumort., Centothecoaceae Ridley, Stipeae Dumort.) and a total of 43 species, half of them in *Aristida* L.

The circumscription of the subfamily has undergone numerous modifications and is accepted here in the sense of Campbell and Davidse and colleagues to include the Centothecoaceae, and in the sense of Barkworth & Everett to include the Stipeae, which were treated as an unplaced tribe by Campbell. Narrower circumscriptions have been proposed by Clayton & Renvoize, Conert (1987), and Renvoize (1982).

In an extensive cluster analysis using 71 characters, Renvoize (1982) demonstrated the importance of leaf-blade anatomy in defining a core group of genera. Several genera, such as *Neyraudia* Hooker f., combine features of the Arundinoideae with those of other traditional subfamilies. In a phylogenetic analysis of the family, Kellogg & Campbell concluded that the subfamily is polyphyletic and probably consists of several groups that are basal to other major clades (i.e., subfamilies) of the grasses.

Danthonia DC. and its allies were included in the Aveneae Dumort. (Pooideae) by Bentham (1883) and Hitchcock (1951) because of their glumes about as long as the spikelets, their several florets per spikelet, and their twisted geniculate lemma awns. Various studies (summarized by Campbell) have pointed out differences between the avenoids and the danthonioids in cytology, leaf anatomy, and embryology. The danthonioids were first accorded tribal status by Zotov, and their similarity to the arundinoids has since been further demonstrated by Hilu & Wright (1982) and Renvoize (1982).

The reed grasses, *Gynerium* Beauv., *Phragmites* Adanson, and *Thysanolaena*

Nees, have chlorenchyma cells with invaginated walls, a bambusoid feature. *Neyraudia* has radiate chlorenchyma (possibly C_4) and is similar and transitional to the Chloridoideae. However, it lacks other chloridoid features such as egg-shaped microhairs, cruciform silica bodies, and triangular subsidiary cells. It has slender arundinoid microhairs, but its embryo is eragrostoid. Its resemblance to *Arundo* and *Phragmites* seems to be due to convergent evolution of habit, not taxonomic affinity (Clayton & Renvoize). The presence of kranz anatomy suggests that species of *Neyraudia* are C_4 , but biochemical and isotopic studies have not yet been conducted to confirm this.

Cortaderia Stapf is a danthonioid genus of about 24 species, with greatest diversity in the Andean and Pampan regions of South America. There are also four species in New Zealand and one in New Guinea. *Cortaderia Selloana* (Schultes) Ascherson & Graebner, pampas grass, is cultivated as an ornamental in warm regions of the United States. Evidence that it spreads from cultivation is lacking, although plants persisting after cultivation have been reported from North Carolina (Radford *et al.*) and Texas (Gould). In California, it is a non-spreading and popular cultivated species. Another cultivated species, *C. jubata* (Lem.) Stapf ("C. atacemensis"), a South American native, is an invasive weed along the California coast (Cowan). In California, only carpellate plants are known and abundant seed is produced apomictically (Costas Lippmann, 1977). It is also a bad weed in New Zealand (Knowles & Ecroyd).

Another danthonioid genus, *Schismus* Beauv., is represented in the southwestern United States and adjacent Mexico by two Eurasian adventives, *S. barbatus* (L.) Thell. (Gould & Moran) and *S. arabicus* Nees.

The Aristideae, the only largely C_4 tribe of the Arundinoideae, include just three genera: *Aristida*, *Sartidia* De Winter, and *Stipagrostis* Nees. It is distinctive both in spikelet morphology and leaf-blade anatomy (Renvoize, 1986). Anatomically, the inner layer of cells in the two-layered bundle sheaths is as large as or larger than the outer layer. The three genera can also be distinguished anatomically. In *Sartidia* the chlorenchyma is not radiate and the photosynthetic pathway is C_3 . *Stipagrostis* (sometimes included in *Aristida*) and *Aristida* are both C_4 genera. In *Stipagrostis* there are no chloroplasts in the inner layer, while in *Aristida* they are present.

The unigeneric tribe Micraireae Pilger is endemic to northern Australia (Clayton & Renvoize; Lazarides). There are 13 species of *Micraira* F. Mueller, all of which grow in very thin soil and show an adaptation to this habitat condition in their ability to recover from extreme desiccation. The genus is unique in the family in having spiral phyllotaxy. The base chromosome number is 10.

The monotypic tribe Thysanolaeneae C. E. Hubb. includes only *Thysanolaena maxima* (Roxb.) O. Kuntze, $2n = 24$, of tropical southern and southeastern Asia. The genus is related to *Phragmites* and may not warrant tribal status (Clayton & Renvoize).

The Centothecaeae are a small group of 11 genera and 26 species (ca. 30, *vide* Clayton & Renvoize; Tenório). Except for *Chasmanthium* Link, which is endemic to southeastern North America, and *Gouldochloa* Valdés, Morden, & Hatch, endemic to northeastern Mexico, they are pantropic. Centothecoids are similar to the herbaceous bambusoids in habit but are decidedly arundinoid

in leaf anatomy. The distinctive embryos have a mesocotyl, lack an epiblast (present in other arundinoids), and have a scutellum cleft and a rolled first leaf. Centothecoid grasses were first accorded taxonomic notice as a subtribe of the Festuceae (= Poaceae) by Bentham (1883). They were subsequently treated as subfam. Centothecoideae (Clayton & Renvoize; Soderstrom; Tenório) but are here considered to be a tribe of the Arundinoideae following Campbell and Davidse and colleagues. A strongly papillate abaxial surface (absent in other grasses) characterizes all genera (Davidse *et al.*). The spikelets are strongly laterally compressed, the caryopsis is laterally flattened, and the hilum is basal and punctiform to shortly linear. Many genera have pulvinate petiolate leaves (*Chasmanthium* and *Gouldochloa* are exceptions), double bundle sheaths, and leaf tissue differentiated into palisade and spongy layers (in grasses otherwise known only in *Lacisias* (Griseb.) Hitchc., Panicoideae A. Br.). Bicellular, often filiform microhairs are commonly present (Davidse *et al.*). A base chromosome number of 12 characterizes nearly all cytologically known genera.

The only large genus, *Zeugites* R. Br., is confined to the New World tropics, as are *Calderonella* Soderstrom & Decker and *Pohlidium* Davidse, Soderstrom, & Ellis. In the Old World tropics are *Bromuniola* Stapf & Hubb., *Centotheca* Desv., *Chevalierella* A. Camus, *Lophatherum* Brongn., and the most primitive genus, *Megastachya* Beauv. Only *Orthoclada* Beauv. has representatives in both the Old World and the New. Tenório proposed (but did not validly publish) a monotypic genus for the African species of *Orthoclada*.

The Stipeae are here included in the Arundinoideae, as has been suggested by Barkworth & Everett and Watson and colleagues. However, Kellogg & Campbell hypothesized that the Stipeae form a clade that is the sister group of the Pooideae. Their amino-acid profiles are intermediate between those of the poooids and the bambusoids. Two-celled microhairs are absent from the Stipeae, as they are from the Pooideae. However, single-celled microhairs, called "pylidial hairs" by Renvoize (1985), are present in some species of *Stipa* and *Oryzopsis* *sensu lato*. These hairs may have evolved from the two-celled, fingerlike microhairs of Bambusoideae. In the Southeast the Stipeae are represented by two genera, *Nassella* (Trin.) Desv. and *Piptochaetium* J. Presl, each of which has been included in *Stipa* by some workers.

The stipoid genus *Oryzopsis* Michx. (including *Piptatherum* Beauv.) is represented by ten or more species in North America, mostly in the West. Three species, *O. asperifolia* Michx., *O. canadensis* (Poiret) Torrey, and *O. racemosa* (J. E. Sm.) Ricker, reach their southern range limits in the mountains of West Virginia and may yet be found in the mountains of North Carolina and Tennessee. The western species *O. hymenoides* (Roemer & Schultes) Ricker, $2n = 48$, comes eastward to Manitoba, Kansas, and western Texas and might be expected as an adventive or waif in Arkansas or Louisiana.

Although the genus *Milium* L. was included in the Stipeae by Clayton & Renvoize, who made the amazing statement that it is merely an awnless version of *Oryzopsis*, it was excluded from the tribe by Barkworth & Everett. According to anatomical data provided by Renvoize (1985), *Milium* is anomalous in the Stipeae in having tapering long cells and in lacking short cells in the leaf epidermis. It is typically poooid in its leaf anatomy, amino-acid profile, susceptibility to rusts, and cytology (see Barkworth & Everett for references).

Haustorial synergids, an unusual micromorphological feature not otherwise known in the grasses, have been found in *Chionochloa* Zotov, *Cortaderia*, *Danthonia* (including *Rytidosperma* Steudel), *Erythranthera* Zotov, *Lamprothyrsus* Pilger, *Pyrrhanthera* Zotov, and *Sieglingia* Bernh. (Philipson & Connor). These massive synergids extend through the micropyle to lie between the ovule and the ovary wall. Normal pyriform synergids, the typical graminoid condition, characterize *Arundo* and *Phragmites*. The ovary in *A. Donax* L. has an unusually long stalk (Bhanwra). The outer epidermal cells of the ovary accumulate an unidentified darkly staining material in *A. Donax* and *P. Karka* (Retz.) Trin. ex Steudel (Bhanwra). Intercalary growth of the chalaza after fertilization characterizes all arundinoid grasses investigated.

Information on the reproductive biology of the arundinoid grasses is incomplete. Most genera have perfect flowers. Species of *Cortaderia* and *Lamprothyrsus* are gynodioecious. The Neotropical *Gynerium sagittatum* (Aublet) Beauv., the only species of the genus, is dioecious. Self-compatibility is known in some species of *Aristida*, *Chasmanthium*, *Chionochloa*, *Cortaderia*, *Danthonia* (including some species assigned to the segregate *Rytidosperma*), *Nassella*, and *Stipa*. *Phragmites australis* is apparently self-incompatible. Apomixis is reported from *Cortaderia jubata* (Philipson, 1978). Four genera in our area (*Aristida*, *Chasmanthium*, *Danthonia*, and *Nassella*) produce cleistogamous as well as chasmogamous spikelets in at least one species. In most genera (all those in the Southeast) the pericarp is fused to the seed coat (ordinary grass caryopsis). However, in several (none in the Southeast) the pericarp is free or separable and the fruit is thus an achene. The genera with achenes are *Amphipogon* R. Br. (Australia), *Anisopogon* R. Br. (Australia), *Dregeochloa* Conert (southern Africa), *Elytrophorus* Beauv. (tropical Africa, India, Australia), *Hakonechloa* Honda (Japan), *Pentameris* Beauv. (southern Africa), *Pyrrhanthera* (New Zealand), and *Urochlaena* Nees (southern Africa). With "reluctantly free pericarp" (Clayton & Renvoize, p. 182) are *Molinia* Schrank (temperate Eurasia) and *Tribolium* Desv. (southern Africa). Except for *Hakonechloa*, all the genera with a free pericarp are confined to the Southern Hemisphere (or to continents derived from Gondwanaland).

The dispersal mechanisms of most arundinoid grasses depend on hairs or awns. The mature spikelets of *Neyraudia* and *Phragmites* break at the joints of the fragile rachilla. The hairs of the rachilla allow the florets to be wind dispersed. The silky, clasping lemmas of *Arundo* probably serve the same function. The long, sharp awns and short, stiff callus hairs of the lemmas of *Aristida* species cause the mature spikelets to stick in fur or skin of animals and thus to be dispersed. Peart, however, examined carcasses of some 100 marsupials and found no awned diaspores in their fur, a surprising result in Australia, where both *Aristida* and *Danthonia* are diverse. African species of *Stipagrostis* have plumose awns, enabling dispersal by wind. The awns of the lemmas of *Danthonia* species are hygroscopic, twisting and coiling and moving the floret into soil crevices, where chances for its germination are improved (Peart). (The hygroscopic awn of *Piptochaetium* may function in the same way.) The long hairs of the lemmas of the Australian *D. tenuior* (Steudel) Conert collapse on drying and rehydrate on wetting, moving the diaspores across soil and into crevices (Peart).

The arundinoid grasses are little known chemically. The leaves of *Phragmites australis* contain flavonoid aglycones (chrysoeriol, isorhamnetin, luteolin, quercetin, tricetin), flavonol and flavone *O*-glycosides (e.g., glucosyl-3 quercetin and glucosyl-7 tricetin), and flavone *C*-glycosides (7,3'-dimethoxy-isoorientin, isoorientin, isoscoparin, swertiajaponin, swertisin). Tricetin and *C*-glycoflavones are known from about 90 percent of the grasses that have been studied (Jay & Viricel). The major flavonoid components of the "flower tissue" (= spikelets?) of *P. australis* are isoswertiajaponin and swertiajaponin (Nawwar *et al.*); also noted were 3'-*O*-gentiobioside and the 3'-*O*-glucoside of swertiajaponin. In addition, there are two flavonol glycosides, rhamnetin 3-*O*-rutoside and rhamnetin 3-*O*-glucoside, a class of compounds rare in grasses. *Phragmites australis* has a minute (0.27 percent) hydrocarbon fraction and no appreciable polyisoprenes (Buchanan *et al.*). The epicuticular layer of *Chionochloa* species contains long-chain carbon compounds (alcohols, aldehydes, alkanes, esters, and fatty acids) up to 52 carbons in length (Savill *et al.*).

Numerous fungi parasitize arundinoid grasses. *Balansia hypoxylon* (Peck) Atk. (anamorph: *Ephelis borealis* Ellis & Everhart) parasitizes *Danthonia compressa* Austin, *D. sericea* Nutt., and *D. spicata* (L.) Beauv., as well as *Aristida glauca* (Nees) Walp. and three species of *Stipa*. It forms segmented sclerotia (Sprague). *Uromyces Danthoniae* McAlp. occurs on six Australian species of *Danthonia*; however, no rusts are reported from any New World species of *Danthonia*. *Puccinia invenusta* H. Sydow & P. Sydow grows on all three species of *Phragmites* in the Old World but has no other hosts. *Puccinia Magnusiana* Körn. (worldwide) and *P. Trabutii* Roum. & Sacc. (northern Africa, southern Asia) both occur on *Arundo* and *Phragmites*. *Puccinia Neyraudiae* H. Sydow & P. Sydow is reported only from *Neyraudia madagascariensis* (Kunth) Hooker f. in southern India (Cummins).

The earliest evidence of arundinoid grasses dates from the Miocene. In a fine study Thomasson (1984) demonstrated anatomical features in silicified remains from Nebraska. Fossilized florets of the extinct stipoid genus *Beriochloa* M. K. Elias were found in the abdominal cavity of a fossilized rhinoceros, *Teleoceras major* Hatcher, and provide direct evidence of grass diet in Miocene times (Voorhies & Thomasson). Quaternary remains of *Phragmites australis* from Egypt could be confidently identified because the internal and external structures of rhizomes and attached stem bases were preserved (El-Saadawi *et al.*).

Plants of the subfamily have some economic importance. None are used for grain. Stems of *Arundo* and *Phragmites* are harvested for paper pulp in Europe and Asia, and they are sometimes important for forage. *Danthonia* species are important for forage in Australia (and to a lesser extent in western North America), as are immature plants of *Aristida* in the American Southwest, but neither is important in these respects in the Southeast. *Nassella leucotricha* is an important cool-weather forage grass in Texas. *Stipa tenacissima* L. is gathered for paper pulp in Algeria and Spain (Hitchcock, 1951).

REFERENCES:

- ACEVEDO DE VARGAS, R. Las especies de Gramineae del género *Cortaderia* en Chile. Bol. Mus. Nac. Hist. Nat. Chile 27: 205-246. 1959.*

- AMARASINGHE, V., & L. WATSON. Comparative ultrastructure of microhairs in grasses. Bot. Jour. Linn. Soc. **98**: 303–319. 1988. [*Amphipogon strictus* R. Br. has “Enneapogon-type” microhairs; cf. RENVOIZE (1986), who found no microhairs in *A. carcinus* F. Mueller.]
- ARBER, A. Monocotyledons. A morphological study. 258 pp. Cambridge, England. 1925. (Reprint, J. Cramer, Weinheim. 1961.)
- . The Gramineae, a study of cereal, bamboo, and grass. 480 pp. New York. 1934.
- BAILEY, L. H., E. Z. BAILEY, & STAFF OF THE BAILEY HORTORIUM. Hortus third. xiv + 1290 pp. New York. 1976.
- BARKWORTH, M. E. Embryological characters and the taxonomy of the Stipeae (Gramineae). Taxon **31**: 233–243. 1982.
- . *Ptilagrostis* in North America and its relationship to other Stipeae (Gramineae). Syst. Bot. **8**: 395–419. 1983. [Includes numerical analysis and many informative illustrations.]
- & J. EVERETT. Evolution in the Stipeae: identification and relationships of its monophyletic taxa. Pp. 251–264 in T. R. SODERSTROM, K. HILU, C. S. CAMPBELL, & M. E. BARKWORTH, eds., Grass systematics and evolution. Washington, D. C. 1987.
- , J. MCNEILL, & J. MAZE. A taxonomic study of *Stipa Nelsonii* (Poaceae) with a key distinguishing it from related taxa in western North America. Canad. Jour. Bot. **57**: 2539–2553. 1979.
- BARTOLOME, J. W. *Stipa pulchra*, a survivor from the pristine prairie. Fremontia **9**(1): 1–6. 1981. [Seed output exceeds 200 kg/ha in dense stands in California.]
- BENTHAM, G. Notes on Gramineae. Jour. Linn. Soc. Bot. **19**: 14–134. 1882.
- . Gramineae. In: G. BENTHAM & J. D. HOOKER, Gen. Pl. **3**: 1074–1215. 1883.
- BHANWARA, R. K. Embryology in relation to systematics of Gramineae. Ann. Bot. II. **62**: 215–233. 1988. [Illustrations.]
- BROWN, W. V. Leaf anatomy in grass systematics. Bot. Gaz. **119**: 170–178. 1958.
- BUCHANAN, R. A., C. L. SWANSON, D. WEISLEDER, & I. M. CULL. Gutta-producing grasses. Phytochemistry **18**: 1069–1071. 1979. [*Phragmites australis*; contains minute hydrocarbon fraction.]
- CAMPBELL, C. S. The subfamilies and tribes of Gramineae (Poaceae) in the southeastern United States. Jour. Arnold Arb. **66**: 123–199. 1985.
- CARO, J. A., & E. SÁNCHEZ. Las especies de *Stipa* (Gramineae) del subgénero *Jarava*. Kurtziana **7**: 61–116. 1973.
- CLAYTON, W. D., & S. A. RENVOIZE. Genera graminum: grasses of the world. Kew Bull. Add. Ser. **13**. [viii +] 389 pp. 1986.
- CLEWELL, A. F. Guide to the vascular plants of the Florida Panhandle. viii + 605 pp. Tallahassee. 1985. [Gramineae, 106–167.]
- CONERT, H. J. Die Systematik und Anatomie der Arundineae. 208 pp. Weinheim, West Germany. 1961.
- . Die *Chionochloa*-Arten von Australien und Neuguinea (Poaceae: Arundinoideae). Senckenberg. Biol. **56**: 153–164. 1975.
- . Current concepts in systematics of the Arundinoideae. Pp. 239–250 in T. R. SODERSTROM, K. HILU, C. S. CAMPBELL, & M. E. BARKWORTH, eds., Grass systematics and evolution. Washington, D. C. 1987.
- & A. M. TÜRPE. Revision der Gattung *Schismus* (Poaceae: Arundinoideae: Danthonieae). Abh. Senckenb. Naturf. Ges. **532**: 1–81. 1974.
- CONNOR, H. E. Reproductive biology in grasses. Pp. 117–132 in T. R. SODERSTROM, K. HILU, C. S. CAMPBELL, & M. E. BARKWORTH, eds., Grass systematics and evolution. Washington, D. C. 1987.
- . Breeding systems in New Zealand grasses. X. Species at risk for conservation. New Zealand Jour. Bot. **26**: 163–167. 1988.
- COSTAS LIPPMANN, M. A. Ecology and reproductive biology of the genus *Cortaderia* in California. 365 pp. Unpubl. Ph.D. thesis, Univ. Calif., Berkeley. 1976.*

- . More on the weedy "pampas grass" in California. *Fremontia* **4**(4): 25–27. 1977. [Good illustrations of *Cortaderia jubata*, *C. Selloana*.]
- COWAN, B. D. The menace of pampas grass. *Fremontia* **4**(2): 14–16. 1976.
- CUMMINS, G. B. The rust fungi of cereals, grasses, and bamboos. xv + 570 pp. New York. 1971.
- DAHLGREN, R. M. T., H. T. CLIFFORD, & P. F. YEO. The families of monocotyledons: structure, evolution, and taxonomy. xii + 520 pp. Berlin, Heidelberg, New York, and Tokyo. 1985.
- DAVIDSE, G., T. R. SODERSTROM, & R. P. ELLIS. *Pohlidium petiolatum* (Poaceae: Centothecaceae), a new genus and species from Panama. *Syst. Bot.* **11**: 131–144. 1986. [Includes much information on circumscription of Centothecaceae.]
- DEDECCA, D. M. Studies on the Californian species of *Stipa* (Gramineae). *Madroño* **12**: 129–160. 1954.
- DEWET, J. M. J. The genus *Danthonia* in grass phylogeny. *Am. Jour. Bot.* **41**: 204–211. 1954.
- . Leaf anatomy and phylogeny in the tribe Danthoneae. *Ibid.* **43**: 175–182. 1956.
- DE WINTER, B. The South African Stipeae and Aristideae. *Bothalia* **8**: 201–404. 1962.
- DORE, W. G., & J. MCNEILL. Grasses of Ontario. *Agr. Canada Monogr.* **26**. 566 pp. 1980.
- ELIAS, M. K. Tertiary prairie grasses and other herbs from the High Plains. *Spec. Pap. Geol. Soc. Am.* **14**: 1–176. 1942. [Includes Stipeae.]
- ELLIS, R. P. Leaf anatomy of the South African Danthoneae (Poaceae). I. The genus *Dregeochloa*. *Bothalia* **12**: 209–213. 1977; XVI. The genus *Urochlaena*. *Ibid.* **18**: 101–104. 1988.
- EL-SAADAWI, W. E., A. A. BADAWI, & A. A. EL-AWAMRI. On silicified rhizome fragments of *Phragmites communis* Trin. from the Pleistocene of El-Fayum, Egypt. *Palaentographica, Abt. B.* **154**: 172–178. pls. 1–5. 1975.
- FOWLER, N. L. Microsite requirements for germination and establishment of three grass species. *Am. Midl. Nat.* **115**: 131–145. 1986. [*Aristida longiseta*, *Stipa leucotricha* (= *Nassella leucotricha*) in Texas.]
- & D. W. DUNLAP. Grassland vegetation of the eastern Edwards Plateau. *Am. Midl. Nat.* **115**: 146–155. 1986. [Ecological information on *Aristida longiseta*, *Stipa leucotricha* (= *Nassella leucotricha*) in Texas.]
- FREITAG, H. The genus *Piptatherum* (Gramineae) in Southwest Asia. *Notes Roy. Bot. Gard. Edinburgh* **33**: 342–408. 1975.
- . The genus *Stipa* (Gramineae) in Southwest and South Asia. *Ibid.* **42**: 355–489. 1985.
- GHOSAL, S., R. K. CHAUDURI, & S. DUTTA. Alkaloids of flowers of *Arundo Donax*. *Phytochemistry* **10**: 2852, 2853. 1971.
- , ———, ———, & S. K. BHATTACHARYA. Occurrence of curarimetic indoles in flowers of *Arundo donax*. *Pl. Med.* **21**: 22–28. 1972. [Nine simple indolic bases isolated.]
- GOULD, F. W. The grasses of Texas. vii + 653 pp. College Station, Texas. 1975.
- & R. MORAN. The grasses of Baja California, Mexico. *San Diego Nat. Hist. Soc. Mém.* **12**. 140 pp. 1981.
- HALL, D. W. The grasses of Florida. 498 pp. Unpubl. Ph.D. thesis, Univ. Florida, Gainesville. 1980.
- HATCH, S. L., J. VALDÉS R., & C. W. MORDEN. *Stipa hirticulmis* (Poaceae: Stipeae), a new species from Nuevo León, Mexico. *Syst. Bot.* **11**: 186–188. 1986.
- HILU, K., & K. WRIGHT. Systematics of Gramineae: a cluster analysis study. *Taxon* **31**: 9–36. 1982.
- HITCHCOCK, A. S. The North American species of *Stipa*. Synopsis of the South American species of *Stipa*. *Contr. U. S. Natl. Herb.* **24**: 215–289. 1925.

- . Manual of the grasses of the United States. ed. 2 (revised by A. CHASE). U. S. Dept. Agr. Misc. Publ. 200. 1054 pp. 1951.
- HOLM, L., J. V. PANCHO, J. P. HERBERGER, & D. L. PLUCKNETT. A geographical atlas of world weeds. (Arabic, Chinese, English, French, German, Hindi, Indonesian, Japanese, Russian, and Spanish introductions.) xlv + 391 pp. New York and other cities. 1979.
- HUBBARD, C. E. Gramineae. Pp. 871–903 in J. HUTCHINSON, The families of flowering plants. Oxford. 1973.
- JACQUES-FÉLIX, H. Les Graminées (Poaceae) d'Afrique tropicale, I: généralités, classification, description des genres. Inst. Recherches Agron. Cult. Vivères Bull. Sci. Paris 8. xi + 345 pp. 1962.*
- JAY, M., & M.-R. VIRICEL. Les flavonoïdes de feuilles du *Phragmites australis*: essai de définition du profil polyphénolique de l'espèce. Phytochemistry 19: 2627, 2628. 1980.
- JOHNSON, B. L. Cytotaxonomic studies in *Oryzopsis*. Bot. Gaz. 107: 1–32. 1945.
- . Natural hybrids between *Oryzopsis* and *Stipa*. I. *Oryzopsis hymenoides* × *Stipa speciosa*. Am. Jour. Bot. 47: 736–742. 1960.
- KAM, Y. K., & J. MAZE. Studies on the relationships and evolution of supraspecific taxa using developmental data. II. Relationships and evolution of *Oryzopsis hymenoides*, *O. virescens*, *O. Kingii*, *O. micrantha*, and *O. asperifolia*. Bot. Gaz. 135: 227–247. 1974.
- KELLOGG, E. A., & C. S. CAMPBELL. Phylogenetic analyses of the Gramineae. Pp. 310–322 in T. R. SODERSTROM, K. HILU, C. S. CAMPBELL, & M. E. BARKWORTH, eds., Grass systematics and evolution. Washington, D. C. 1987.
- KNOWLES, B., & C. ECROYD. Species of *Cortaderia* (pampas grass and toetoe) in New Zealand. Rotorua Forest Res. Inst. Bull. 105: ii + 24 pp. 1985. [Adventive *C. jubata* a forest weed.]
- LAZARIDES, M. *Micraira*. Brunonia 2: 67–84. 1979. [Australian endemic genus forms tribe Micraireae.]
- LONG, R. W., & O. LAKELA. A flora of tropical Florida. xvii + 962 pp. Coral Gables, Florida. 1971. [Gramineae, 132–202.]
- MATTHEI, O. R. Estudio crítico de las gramíneas del género *Stipa* en Chile. Gayana Bot. 13: 1–137. 1965. [Thirty-nine species; *Nassella* excluded; illustrations.]
- MAZE, J. Notes on the awn anatomy of *Stipa* and *Oryzopsis* (Gramineae). Syesis 5: 169–171. 1972.
- & L. R. BOHM. Comparative embryology of *Stipa Elmeri* (Gramineae). Canad. Jour. Bot. 51: 235–247. 1973.
- & S. C. LIN. A study of the mature megametophyte of *Stipa Elmeri*. Canad. Jour. Bot. 53: 2958–2977. 1975.
- MUNZ, P. A. A California flora. 1681 pp. Berkeley & Los Angeles. 1959.
- NAWWAR, M. A. M., H. I. EL SISSI, & H. H. BARACAT. The flavonoids of *Phragmites australis* flowers. Phytochemistry 19: 1854–1856. 1980.
- PALMER, P. G., & A. E. TUCKER. A scanning electron microscope survey of the epidermis of East African grasses, I. Smithson. Contr. Bot. 49: 1–84. 1981. [*Asphenatherum glaucum*, *Bromuniola Gossweileri*, *Crinipes abyssinicus*, *Elytrophorus globularis*, *Haemobrochia Bullockii*, *Megastachya mucronata*, *Neyraudia arundinacea*, *Orthocladia africana*, *Pentastichis borussica*, *Phragmites mauritianus*, *Triraphis Schinzii*.]
- PARODI, L. R. Las especies de *Stipa* del subgénero *Pappostipa* de la Argentina y Chile. Revista Argent. Agron. 27: 65–106. 1960.
- PEART, M. H. Experiments on the biological significance of the morphology of seed-dispersal units in grasses. Jour. Ecol. 67: 843–863. 1979.
- PHILIPSON, M. N. Haustorial synergids in *Cortaderia* (Gramineae). New Zealand Jour. Bot. 15: 777, 778. 1977.
- . Apomixis in *Cortaderia jubata* (Gramineae). Ibid. 16: 45–59. 1978.

- . The haustorial synergids of *Cortaderia* (Gramineae) at maturity. *Acta Soc. Bot. Polon.* **50**: 151–160. 1981.*
- & H. E. CONNOR. Haustorial synergids in danthonioid grasses. *Bot. Gaz.* **145**: 78–82. 1984.
- RADFORD, A. E., H. E. AHLES, & C. R. BELL. Manual of the vascular flora of the Carolinas. lxi + 1183 pp. Chapel Hill. 1968.
- REEDER, J. R. The embryo in grass systematics. *Am. Jour. Bot.* **44**: 756–768. 1957.
- RENVOIZE, S. A. The subfamily Arundinoideae and its position in relation to a general classification of the Gramineae. *Kew Bull.* **36**: 85–102. 1982.
- . A survey of the leaf blade anatomy in grasses. VI. Stipeae. *Ibid.* **40**: 731–736. 1985; VIII. Arundinoideae. *Ibid.* **41**: 323–342. 1986.
- ROBINSON, E. R. Naturalized species of *Cortaderia* (Poaceae) in southern Africa. *S. Afr. Jour. Bot.* **3**: 343–346. 1984. [Adventives *C. Selloana*, *C. jubata* are invasive weeds.]
- ROSENGURTT, B., & B. R. ARRILLAGA DE MAFFEI. Nuevas especies y sinopsis de *Stipa* en el Uruguay. *Bol. Fac. Agron. Montevideo* **72**: 1–34. 1964.
- SAVILL, M. G., R. BICKESTAFF, & H. E. CONNOR. Interspecific variation in epicuticular waxes of *Chionochloa*. *Phytochemistry* **27**: 3499–3507. 1988. [Each species has distinct profile of long-chain carbon compounds.]
- SCHECHTER, Y., & B. L. JOHNSON. A new species of *Oryzopsis* (Gramineae) from Wyoming. *Brittonia* **18**: 342–347. 1966.
- & ———. The probable origin of *Oryzopsis contracta*. *Am. Jour. Bot.* **55**: 611–618. 1968.
- SODERSTROM, T. R. The grass subfamily Centostecoideae. *Taxon* **30**: 614–616. 1981. [Validation of subfamily: synopsis of included genera.]
- & H. F. DECKER. *Calderonella*, a new genus of grasses and its relationships to the centostecoid genera. *Ann. Missouri Bot. Gard.* **60**: 427–441. 1973.
- SPELLENBERG, R. W., & L. E. MEHLENBACHER. Anatomical and cytological studies of an intergeneric hybrid, *Oryzopsis Hendersonii* × *Stipa Lemmonii* (Gramineae). *Canad. Jour. Bot.* **49**: 1565–1574. 1971.
- SPRAGUE, R. Diseases of cereals and grasses in North America (fungi, except rusts and smuts). xvi + 538 pp. New York. 1950.
- STAPF, O. The pampas grasses. *Fl. Sylva* **3**: 171–176. 1905. [*Cortaderia*.]
- TAIRA, H. Studies on amino acid contents in plant seeds IV. Amino acid contained in the seed of Gramineae (part 3). *Bot. Mag. Tokyo* **79**: 36–48. 1966.
- TENÓRIO, E. C. The subfamily Centostecoideae (Gramineae). xii + 420 pp. Unpubl. Ph.D. dissertation, Univ. Maryland, College Park. 1978.
- THOMASSON, J. R. Late Cenozoic grasses and other angiosperms from Kansas, Nebraska, and Colorado: biostratigraphy and relationships to living taxa. *Kansas Geol. Surv. Bull.* **218**: 1–68. 1979.
- . *Paleoeriucoma* (Gramineae, Stipeae) from the Miocene of Nebraska: taxonomic and phylogenetic significance. *Syst. Bot.* **5**: 233–240. 1980 [1981].
- . Miocene grass (Gramineae: Arundinoideae) showing external micromorphological and internal anatomical features. *Bot. Gaz.* **145**: 204–209. 1984.
- TOMLINSON, K. L. Comparative anatomical studies in *Danthonia* sensu lato (Danthoniaceae: Poaceae). *Aliso* **11**: 97–114. 1985. [Leaf and lodicule anatomy.]
- TOWNROW, J. E. S. The genus *Stipa* in Tasmania. Part 3—revised taxonomy. *Pap. Proc. Roy. Soc. Tasmania* **112**: 227–287. 1978. [Twelve species described and illustrated; all indigenous.]
- TSVELEV, N. N. *Zlaki SSSR*. 2 parts. Leningrad. 1976. (Grasses of the Soviet Union. English translation by B. R. Sharma. 2 parts (1196 pp.). Rotterdam. 1984.) [See pp. 848–936 in transl.]
- TUTIN, T. G. Arundineae, Cortaderieae, Danthonieae, Molinieae, and Aristideae (Gramineae). Pp. 252–255 in T. G. TUTIN *et al.*, eds., *Flora Europaea*. Vol. 5. Cambridge, England. 1981.

- VALDÉS R., J., C. W. MORDEN, & S. L. HATCH. *Gouldochloa*, a new genus of centothecoid grasses from Tamaulipas, Mexico. *Syst. Bot.* **11**: 112–119. 1986. [Illustrations.]
- VICKERY, J. W., S. W. L. JACOBS, & J. EVERETT. Taxonomic studies in *Stipa* in Australia. *Telopea* **3**: 1–133. 1986.
- VOORHIES, M. R., & J. R. THOMASSON. Fossil grass anthoecia within Miocene rhinoceros skeletons: diet in an extinct species. *Science* **206**: 331–333. 1979.
- WALTERS, S. M., *et al.*, eds. The European garden flora. Vol. 2. 318 pp. Cambridge, England, and other cities. 1984. [Arundinoideae, 34–36; six genera.]
- WATSON, L., H. T. CLIFFORD, & M. J. DALLWITZ. The classification of the Poaceae: subfamilies and supertribes. *Austral. Jour. Bot.* **33**: 433–484. 1985.
- WUNDERLIN, R. P. Guide to the vascular plants of Central Florida. viii + 472 pp. Tampa and other cities. 1982. [Poaceae, 51–96.]
- YATES, H. O. Morphology and cytology of *Uniola* (Gramineae). *Southwest. Nat.* **11**: 145–189. 1966.
- . Revision of grasses traditionally referred to *Uniola*, II. *Chasmanthium*. *Ibid.* 415–455. 1966 [1967].
- YEOH, H. H., & L. WATSON. Systematic variation in amino acid compositions of grass caryopses. *Phytochemistry* **20**: 1041–1051. 1981.
- ZOTOV, V. D. Synopsis of the grass subfamily Arundinoideae in New Zealand. *New Zealand Jour. Bot.* **1**: 78–136. 1963.

KEY TO THE GENERA OF ARUNDINOIDEAE IN THE SOUTHEASTERN UNITED STATES

General characters: *perennial or annual herbaceous plants of dry to wet places. Leaf blades linear to lanceolate, flat or involute. Inflorescences paniculate (rarely racemose), terminal and/or axillary. Spikelets laterally compressed or subterete. Glumes 2, conspicuous; flowers 1 to several, disarticulating above glumes; lemmas often awned.*

- A. Flowers imperfect or perfect (plants gynodioecious), forming dense tussocks. [Cortaderia.]
- A. Flowers perfect (lower florets sometimes imperfect); plants rhizomatous, or loosely to densely caespitose.
- B. Plants large; leaves cauline only.
- C. Rachillas glabrous; lemmas pilose throughout. 1. *Arundo*.
- C. Rachillas pilose; lemmas glabrous or pilose on margins only.
- D. External ligules a line of hairs; lateral inflorescences often present; lemmas with excurved awn. 3. *Neyraudia*.
- D. External ligules absent; inflorescences strictly terminal; lemmas awnless. 2. *Phragmites*.
- B. Plants small to medium; leaves basal and cauline.
- E. Spikelets subterete, 1-flowered; lemmas indurate, involute.
- F. Lemma awns 1-parted.
- G. Paleas rudimentary. 6. *Nassella*.
- G. Paleas slightly longer than and protruding from apex of lemmas. 7. *Piptochaetium*.
- F. Lemma awns 3-parted. 5. *Aristida*.
- E. Spikelets laterally flattened, 2- to 12-flowered; lemmas membranaceous, conduplicate.
- H. Glumes and spikelets about the same length; lemmas awned; caryopses subcylindrical; leaves mostly basal. 4. *Danthonia*.
- H. Glumes much shorter than spikelets; lemmas awnless; caryopses laterally flattened; leaves mostly cauline. 8. *Chasmanthium*.

Tribe ARUNDINEAE Dumortier, Obs. Gram. Belg. 82. 1824.

1. **Arundo** Linnaeus, Sp. Pl. 1: 81. 1753; Gen. Pl. ed. 5. 35. 1754.

Very large perennials of sunny, damp soils and shallow, fresh waters. Roots fibrous; rhizomes stout. Stems stout, solid, glabrous; nodes glabrous. Leaves several, cauline; sheath glabrous; ligule membranaceous, margin minutely ciliate; blade lanceolate, auriculate, the margin and apex scabridulous; cross veins numerous. Inflorescences single, terminal, much branched, plumose. Spikelets lanceolate; glumes 2, lanceolate, as long as the spikelets; rachilla slender, flattened, glabrous; florets [1 or] 2; lemmas lanceolate, clasping, membranaceous to firm, 5- to 9-nerved, the lower $\frac{1}{2}$ pilose; paleas lanceolate, $\frac{1}{2}$ – $\frac{2}{3}$ as long as the lemmas, hyaline, the margin ciliate. Stamens 3; anthers linear, the apex and base of thecae divergent. Ovaries slenderly cylindrical; styles 2, free, very slender; stigmas slightly longer than styles, plumose, laterally exserted. Caryopses oblong, smooth; hilum short; embryo large. Base chromosome number 12. LECTOTYPE SPECIES: *A. Donax* L.⁴ (Name from Latin word for cane.)—GIANT REED.

A genus of three species native to the Old World. *Arundo Donax* L., $2n = 110$, the only wide-ranging species, occurs from Spain to India. The other species are *A. Plinii* Turra, $2n = 72$, of the Mediterranean region, and *A. formosana* Hackel, endemic to Taiwan.

Arundo Donax is naturalized from seed of planted specimens in the southeastern and southwestern United States. I have examined specimens from Virginia, North Carolina, Florida, Louisiana, Texas, and California.

Seed set is reported to be poor in *Arundo Donax* in India because of the failure of meiosis in a majority of the ovules (Bhanwra).

The reeds for woodwind instruments are cut from the stems of *Arundo Donax* (Clayton & Renvoize). The stems are used for thatch in southern Europe, Asia, and Africa, cellulose for paper pulp, and fodder. *Arundo Donax* is also cultivated around pools and on streambanks as an ornamental.

REFERENCES:

Under subfamily references see BAILEY *et al.*; BHANWRA; CLAYTON & RENVOIZE; CONERT (1961); MUNZ; RENVOIZE (1986); TUTIN; and WALTERS *et al.*

2. **Phragmites** Adanson, Fam. Pl. 2: 34, 559. 1763.

Tall, rhizomatous perennials of swamps, marshes, ditches, and roadsides. Roots fibrous; rhizomes stout, scaly, horizontal to oblique; long stolons often produced by terrestrial plants. Stems glabrous, often glaucous. Leaves cauline, 10–15; sheaths overlapping, with margin free for entire length, glabrous; ligule

⁴Linnaeus included six species in his diverse genus *Arundo*. The genus was typified by removal of five of these: *A. Bambos* L. to *Bambusa* ("Bambus") by Gmelin (Syst. Nat. ed. 13 2(1): 579. 1791); *A. Phragmites* L. to *Phragmites* by Adanson (Fam. Pl. 34, 559. 1763); *A. epigejos* L. and *A. Calamagrostis* L. to *Calamagrostis* by Adanson (*Ibid.* 31, 530); and *A. arenaria* L. to *Ammophila* by Host (Gram. Austr. 4: 24. pl. 41. 1809), leaving only *A. Donax*. Linnaeus (Gen. Pl. 35. 1754) described the lemmas as pilose (valvulae . . . basi lanugo") in *Arundo*, a feature fitting only *A. Donax* among the species he included in *Arundo*.




FIGURE 1. Tribe Arundineae, spikelets or their parts. a, b, *Arundo Donax*: a, spikelet, $\times 3$; b, floret (palea and lemma enclosing flower) from adaxial side (note glabrous rachilla segment and hairs along edge of lemma), $\times 3$. c, d, *Phragmites australis*: c, spikelet (parts spread out and hairs omitted for clarity), $\times 4$; d, floret from adaxial side (note abundant hairs from rachilla segment), $\times 4$. e, f, *Danthonia spicata*: e, spikelet, $\times 3$; f, floret from adaxial side (note geniculate awn borne from middle of notch between bilobed lemma apex), $\times 5$.

a short membrane bearing dense fringe of multicellular hairs; blades distichous below, becoming secund above, linear-lanceolate, constricted basally, widest at about $\frac{1}{3}$ the length, tapered to a long, attenuate tip, flat, the midvein much wider and thicker than the lateral veins, cross veinlets visible on the abaxial surface. Inflorescences terminal, solitary, plumose, each node bearing several primary branches (1 or 2 of these larger than the others); nodes of the primary branches each with 1 or 2 secondary branches; secondary branches bearing single tertiary branch at each node; inflorescence axis, secondary, and tertiary branches terminated by solitary spikelets; nodes of inflorescence with ring of long, multicellular trichomes (these giving a woolly appearance to inflorescence). Spikelets numerous, lanceolate; glumes 2, unequal, the second shorter than the florets; rachilla pilose, giving the inflorescence a silky or fluffy appearance; flowers 2 to several, the lowest staminate or empty, the succeeding perfect; lemmas lanceolate, long-acuminate, 3-nerved; paleas less than $\frac{1}{2}$ as long as the lemmas. Stamens 3 (lacking, 1, or 2 in lowest, staminate flower); anthers ellipsoid. Ovaries oblong; styles separate, short; stigmas plumose. Caryopses subterete, oblong; styles persistent, indurate; basal furrow broad; hilum $\frac{1}{3}$ as long as grain. Base chromosome number 12. TYPE SPECIES: *Arundo Phragmites* L. = *P. australis* (Cav.) Trin. ex Steudel. (Name from Greek *phragmites*,

hedge, probably applied to this genus because of its dense, hedgelike growth along ditches and waterways.)—COMMON REED.

A genus of three (or more) species, nearly cosmopolitan in distribution. The only species in the New World, *Phragmites australis* (*P. communis* Trin.), is widespread, but not especially abundant or common, in the Southeast.

Occurring on all continents except Antarctica, *Phragmites australis* may have the widest distribution of any angiosperm. It was present in the American Southwest at least 1000 years before European contact (Kane & Gross), and in southern New England at least 3000 years ago (Niering & Warren). Reported in 1843 as occasional throughout New York State (Torrey), it was not reported from Georgia or South Carolina by Elliot (1821–1824) and remained unknown in those states until the 1970's (Stalter). It is now abundant in New York, where it may be in part adventive (Mitchell). It spread rapidly on Long Island during the early twentieth century, which may be explained by the introduction of a more aggressive biotype from Europe or elsewhere. Distinctive morphotypes, presumably genetically based, have been reported from southern Louisiana (Schott & White). The complex and confused taxonomy of Old World populations makes it difficult to match the North American populations with any of the numerous European varieties and cultivated forms.

Phragmites japonicus Steudel, $2n = 48$, was recognized by Tsvelev, who distinguished it from *P. australis* by its shorter glumes and spikelets and its zigzag (not straight), elongate rhizomes. The species occurs in Japan, China, and the extreme eastern Soviet Union. *Phragmites Karka* (Retz.) Trin., $2n = 18, 36, 38, 48$, grows in tropical Africa, southeastern Asia, and northern Australia (where its distribution complements that of the more temperate *P. australis*). *Phragmites mauritianus* Kunth is found in tropical Africa and the islands of the Indian Ocean (Fanshawe). The species are barely distinguishable (Clayton & Renvoize).

There is considerable variation in the chromosome complement of *Phragmites australis*. Haslam (1972) reported diploid numbers of 36, 48, 84, and 96. Aneuploids are also reported: $2n = 42, 44, 46, 49, 50, 51, 52, 54, 56$ (Gorenflot *et al.*, 1972). In Europe tetraploids and octaploids account for most populations; hexaploids are rare, being reported from Sweden (Björk), the Mediterranean region, and Iran and Afghanistan (Bahrman & Gorenflot). In the New World hexaploids have been reported from Costa Rica (Pohl). Tetraploid and octaploid clones grow together in the Danube Delta (Bahrman & Gorenflot). No correlation between ploidy, habitat, and geography was found in a survey of 40 European populations (Raicu *et al.*). Apparently the only report from North America is $2n = 48$ from a Canadian population (Hunter).

Bahrman & Gorenflot reported variation in soluble proteins of leaves in an extensive survey of European, North African, and Central Asian populations. Principal-components analysis revealed four major groups on the basis of isozymes present. The diversity of isozymes could not be correlated with ploidal level. The greatest diversity in isozymes was found in populations in Iran and Afghanistan, while the greatest variation in chromosome number was in southern Europe. Evidently, macromolecular diversification in this species has proceeded independently from chromosomal changes.

There are numerous reports worldwide of lack of seed set in *Phragmites australis*. A complete explanation has yet to be provided, and the question offers possibilities for further research. Meiosis is regular in certain populations from France (Cartier & Lenoir). However, chromosome fragmentation during microsporogenesis was detected in populations from Ireland that failed to set seed (Curran). In populations from India, megasporogenesis was normal, but microsporogenesis was arrested at the tetrad stage, and there was no mitotic division to produce the vegetative and generative nuclei (Satyamurty & Seshavatharam). Abundant seed set has been reported in populations from Afghanistan and Iran (Bahrman & Gorenflot), South Africa (Curran), Minnesota (Harris & Marshall), and the Mackenzie District, Northwest Territories, Canada (Cody). In Swedish populations, however, reproduction by seed was limited: only 0.3–8.0 percent of the florets produced viable seed (Gustafsson & Simak). No viable fruits have been reported from Ontario populations, and reproduction there has apparently been entirely vegetative (Dore & McNeill). Self-incompatibility may account for low seed set because clonal growth has produced genetically homogeneous populations (genets).

Germination is influenced by temperature, with the rate increasing linearly from 16 to 26°C, while the number of days needed for germination decreases from 25 at 16°C to only ten at 26°C. The seeds cannot germinate under more than 5 cm of water. Seed germination is little affected by salt concentrations below one percent, reaching a limit of tolerance at two percent (Kim *et al.*) in Korean populations. Plants of *Phragmites australis* are able to grow in brackish as well as fresh water. Evidently tolerance of salinity varies widely (Hocking *et al.*): a maximum tolerance for mature plants was reported as 1.2 percent in Britain, 2.9 percent in New York, and 4.0 percent on the Red Sea coast.

For detailed reviews of the autecology and physiological ecology of *Phragmites australis*, see Haslam (1972, 1973) and Hocking and colleagues. For a popular nontechnical account of general ecology and economic importance, see Brown.

Plants of *Phragmites* are used by humans in several ways. Stems are used for basketry and thatch in several European countries, and as a source of pulp for paper production, especially in eastern Europe. Immature stems provide forage for cattle in Australia (Hocking *et al.*) and southern Africa. *Phragmites australis* is an invasive weed in the northeastern United States. The rhizomes are able to grow under pavement and cause damage by cracking and piercing it (Amano & Maki; Hocking *et al.*). Unwanted populations can be effectively removed by cutting them several times annually, particularly in midsummer when rhizome starch reserves are lowest. Dense stands provide wildlife habitat in Europe (Bibby & Lunn). Reeds are planted on recently drained lands in the Netherlands and Japan (Kamio) to remove excess water and prepare the soil for agriculture.

REFERENCES:

- Under subfamily references see CLAYTON & RENVOIZE; DORE & MCNEILL; HOLM *et al.*, MUNZ; and TSVELEV.
- AMANO, K., & T. MAKI. Studies on the method of random paving for the slope protection of embankments. (Part 3) The mechanism of the destruction in asphalt concrete

- paving by reeds (*Phragmites communis* Trinius). (In Japanese; English summary.) Jour. Agr. Sci. Tokyo **29**: 89–100. 1984.
- APINIS, A. E., C. G. C. CHESTERS, & H. K. TALIGOOLA. Colonisation of *Phragmites communis* leaves by fungi. Nova Hedwigia **23**: 113–124. 1972.
- , ———, & ———. Microfungi colonizing nodes and internodes of aerial standing culms of *Phragmites communis* Trin. *Ibid.* **26**: 495–507. 1975.
- BAHRMAN, N., & R. GORENFIOT. Apport des protéines foliaires solubles dans l'interprétation du complexe polyploïde du *Phragmites australis* (Cav.) Trin. ex Steudel. Rév. Gén. Bot. **90**: 177–184. 1983.
- BAYLY, I. L., & T. A. O'NEILL. Seasonal ionic fluctuations in a *Phragmites communis* community. Canad. Jour. Bot. **50**: 2103–2109. 1972.
- BEST, E. P. H., M. ZIPPIN, & J. H. A. DASSEN. Growth and production of *Phragmites australis* in Lake Vechten (The Netherlands). Hydrobiol. Bull. **15**: 165–173. 1981.
- BIBBY, C. J., & J. LUNN. Conservation of reed beds and their avifauna in England and Wales. Biol. Conserv. **23**: 167–186. 1982. [*Phragmites australis*.]
- BJÖRK, S. Ecologic investigations of *Phragmites communis*. Studies in theoretic and applied limnology. Folia Limnol. Scand. **14**: 3–248. 1967.
- BORNKAMM, R., & F. RAGHI-ATRI. Über die Wirkung unterschiedlicher Gaben von Stickstoff und Phosphor auf die Entwicklung von *Phragmites australis* (Cav.) Trin. ex Steudel. Arch. Hydrobiol. **105**: 423–441. 1986. [Effects of nitrogen and phosphorus on growth.]
- BRIX, H. Light dependent variations in the composition of the internal atmosphere of *Phragmites australis* (Cav.) Trin. ex Steudel. Aquatic Bot. **30**: 319–329. 1988.
- BROWN, L. Reed discovery. Horticulture **59**(2): 32–37. 1981.
- CARTIER, D., & A. LENOIR. Contribution à l'étude du développement de l'ovule du *Phragmites australis* (Cav.) Trin. ex Steud. Rev. Gén. Bot. **87**: 289–295. 1980.
- CHASHCHUKHIN, V. A. Ecological aspects of the gas regime in the rhizome of the common reed. Soviet Jour. Bot. **10**: 68, 69. 1979.
- CLAYTON, W. D. Studies in the Gramineae: XIV. Kew Bull. **21**: 111–117. 1967. [World-wide synopsis of *Phragmites*; three species.]
- . The correct name of the common reed. Taxon **17**: 168, 169. 1968.
- CODY, W. J. A contribution to the knowledge of the flora of the southwestern Mackenzie District, N. W. T. Canad. Field-Nat. **77**: 108–120. 1963.
- CURRAN, P. L. Fertility of *Phragmites communis*. Irish Nat. Jour. **16**: 242. 1969.
- DAVIS, A. N., & T. L. BRIGGS. Dispersion pattern of aerial shoots of the common marsh reed *Phragmites australis* (Poaceae). Rhodora **88**: 325–330. 1986.
- DE LA CRUZ, A. A. The production of pulp from marsh grass. Econ. Bot. **32**: 46–50. 1978.
- DENNY, P. Permanent swamp vegetation of the Upper Nile. Hydrobiologia **110**: 79–90. 1984. [*Phragmites Karka*.]
- DINKA, M. The effect of mineral nutrient enrichment of Lake Balaton on the common reed (*Phragmites communis*). Folia Geobot. Phytotax. **21**: 65–84. 1986.
- DRUM, R. W. Electron microscopy of opaline phytoliths in *Phragmites* and other Gramineae. (Abstract.) Am. Jour. Bot. **55**: 713. 1968.
- DURSKA, B. Changes in the reed (*Phragmites communis* Trin.) condition caused by diseases of fungal and animal origin. Pol. Arch. Hydrobiol. **17**: 373–396. 1970.*
- DYKJOVÁ, D. Ecomorphoses and ecotypes of *Phragmites communis* Trin. (In Czech; English summary.) Preslia **43**: 120–138. 1971.
- & D. HRADECKÁ. Production ecology of *Phragmites communis*. 1. Relations of two ecotypes to the microclimate and nutrient conditions of habitat. Folia Geobot. Phytotax. **1**: 23–61. 1966.
- , K. VÉBER, & K. PRIBÁN. Productivity and root/shoot ratio of reedswamp species growing in outdoor hydroponic cultures. Folia Geobot. Phytotax. **6**: 233–254. 1971.
- ELLIOT, S. Sketch of the botany of South Carolina and Georgia. 2 vols. 606 + 743 pp. Charleston, South Carolina. 1821–1824.

- FANSHAWE, D. B. The biology of the reed—*Phragmites mauritianus* Kunth. *Kirkia* **8**: 147–150. 1972.
- FERNALD, M. L. The generic name *Phragmites*. *Rhodora* **24**: 55, 56. 1922.
- . *Phragmites communis* Trin. var. *Berlandieri*. *Ibid.* **34**: 211, 212. 1932.
- . *Phragmites communis* vs. *P. maximus*. *Ibid.* **43**: 286, 287. 1941.
- GALLAGHER, J. L., & M. J. MILLS. Seasonal patterns in storage carbohydrates in *Phragmites australis*. (Abstract.) *Am. Jour. Bot.* **73**: 667. 1986.
- GORENFLOT, R. Le complexe polyploïde du *Phragmites australis* (Cav.) Trin. ex Steud. (= *P. communis* Trin.). *Bull. Soc. Bot. France* **123**: 261–271. 1976.
- , J.-M. HUBAC, M. JAY, & P. LA LANDE. Geographic distribution, polyploidy, and patterns of flavonoids in *Phragmites australis* (Cav.) Trin. ex Steud. *Proc. NATO Adv. Stud. Inst. G.* **1**: 474–478. 1982.*
- , P. RAICU, D. CARTIER, I. CIOBANU, V. STOIAN, & S. STAIKU. Polyploid complex of *Phragmites communis* Trin. *Compt. Rend. Acad. Sci. Paris, D.* **274**: 1501–1504. 1972.
- & M. SANEI CHARIAT-PANAHI. Le complexe polyploïde du *Phragmites australis* (Cav.) Trin. ex Steud. (= *P. communis* Trin.) en Iran. *Rév. Cytol. Biol. Vég. Bot.* **2**: 67–81. 1979.
- GORHAM, E., & W. H. PEARSALL. Production ecology III. Shoot production in *Phragmites* in relation to habitat. *Oikos* **7**: 206–214. 1956.
- GUSTAFSSON, Å., & M. SIMAK. X-ray photography and seed sterility in *Phragmites communis* Trin. *Hereditas* **49**: 442–450. 1963.
- HALDEMANN, C., & R. BRÄNDLE. Seasonal variations of reserves and of fermentation processes in wetland plant rhizomes at the natural site. *Flora* **178**: 307–313. 1986.
- HANSSON, L.-Å., & W. GRANÉLI. Effects of winter harvest on water and sediment chemistry in a stand of reed (*Phragmites australis*). *Hydrobiologia* **112**: 131–136. 1984.
- HARRIS, S. W., & W. H. MARSHALL. Experimental germination of seed and establishment of seedlings of *Phragmites communis*. *Ecology* **41**: 395. 1960.
- HASLAM, S. M. Stem types of *Phragmites communis* Trin. *Ann. Bot. II.* **33**: 127–131. 1969a.
- . The development and emergence of buds in *Phragmites communis* Trin. *Ibid.* **289**–301. 1969b.
- . The development of shoots in *Phragmites communis* Trin. *Ibid.* **695**–709. 1969c.
- . The development of the annual population in *Phragmites communis* Trin. *Ibid.* **34**: 571–591. 1970.
- . Biological flora of the British Isles. *Phragmites communis* Trin. *Jour. Ecol.* **60**: 585–610. 1972.
- . Some aspects of the life history and autecology of *Phragmites communis* Trin., a review. *Polak. Arch. Hydrobiol.* **20**: 79–100. 1973.
- HO, Y. B. Shoot development and production studies of *Phragmites australis* (Cav.) Trin. ex Steudel in Scottish lochs. *Hydrobiologia* **64**: 215–222. 1979.
- HOCKING, P. J., C. M. FINLAYSON, & A. J. CHICK. The biology of Australian weeds. 12. *Phragmites australis* (Cav.) Trin. ex Steud. *Jour. Austral. Inst. Agr. Sci.* **1983**: 123–132. 1983. [Detailed review emphasizing ecology.]
- HU, Y. H., & C. L. LI. Comparative anatomy of the culm and fibers of *Phragmites communis* and *Miscanthus sinensis*. (In Chinese; English summary.) *Acta Bot. Sinica* **11**: 252–260. 1963.
- HUNTER, A. W. S. A karyosystematic investigation in the Gramineae. *Canad. Jour. Res.* **11**: 213–241. 1934.
- HURGHISIU, I. Soluble proteins in *Phragmites australis* (Cav.) Trin. ex Steud. growing in different ecological conditions. *Hydrobiologia* **15**: 275–278. 1977.
- INGRAM, H. A. P., A. M. BARCLAY, A. M. COUPAR, J. G. GLOVER, B. M. LYNCH, & J. I. SPRENT. *Phragmites* performance in reed beds in the Tay Estuary. *Proc. Roy. Soc. Edinburgh* **78B**: 89–107. 1980.
- KAMIO, A. Studies on the drying of marshy and heavy clay soil ground by means of

- vegetations—on the process of polder land drainage and structural changes of *Phragmites communis* community in the Hachirogata central polder. Jap. Jour. Ecol. **32**: 357–364. 1982.
- KANE, A. E., & G. T. GROSS. Anasazi communities at Dolores: early Anasazi sites in the Sagehen Flats area. 985 pp. Denver. 1986. [*Phragmites* plant parts associated with several ruins in southwestern Colorado dated A.D. 600–800.]
- KANETA, M., & N. SUGIYAMA. The constituents of *Arthraxon hispidus* Makino, *Miscanthus tinctorius* Hackel, *Miscanthus sinensis* Anderss., and *Phragmites communis* Trinius. Bull. Chem. Soc. Japan **45**: 528–531. 1972.
- KAUPPI, P., J. SELKÄINAHÖ, & P. PUTTONEN. A method for estimating above-ground biomass in *Phragmites* stands. Ann. Bot. Fenn. **20**: 51–55. 1983.
- KIM, K. S., Y. S. MOON, & C. K. LIM. Effect of NaCl on germination of *Atriplex Gmelinii* and *Phragmites communis*. (In Korean; English abstract.) Korean Jour. Bot. **28**: 253–259. 1985.
- LAU, E., M. GOLDOFTAS, V. D. BALDWIN, P. DAYANANDAN, J. SRINIVASAN, & P. B. KAUFMAN. Structure and localization of silica in the leaf and internodal epidermal system of the marsh grass *Phragmites australis*. Canad. Jour. Bot. **56**: 1696–1701. 1978.
- LAWRENCE, D. B. Arboretum's reed marsh in historical perspective: a plea for conservation of a natural resource. Minn. Agr. Exper. Sta. Misc. Rep. **111**: 24–27. 1972.
- LEFOR, M. W. The common reed, *Phragmites australis* (Cav.) Trin. ex Steud. (Gramineae): a brief overview. Newsletter Conn. Bot. Soc. **15**(1): 7–10. 1987.
- LENOIR, A., V. STOIAN, D. CARTIER, R. GORENLOT, & P. RAICU. Polyplöidie et méiose pollinique du *Phragmites australis* (Cav.) Trin. ex Steud. Compt. Rend. Acad. Sci. Paris, D. **280**: 621–624. 1975.
- LUTHER, H. Beobachtungen über die fruktifikative Vermehrung von *Phragmites communis* Trin. Acta Bot. Fenn. **46**: 1–18. 1950.
- MANTLE, P. G. Studies on *Claviceps purpurea* (Fr.) Tul. parasitic on *Phragmites communis* Trin. Ann. Appl. Biol. **63**: 425–434. 1969.
- METZLER, K., & R. ROZSA. Additional notes on the tidal wetlands of the Connecticut River. Newsletter Conn. Bot. Soc. **15**(1): 1–6. 1987.
- MITCHELL, R. S. A checklist of New York State plants. New York State Mus. Bull. **458**. x + 272 pp. 1986.
- MOOK, J. H., & J. VAN DER TOORN. The influence of environmental factors and management on stands of *Phragmites australis*. II. Effects on yield and its relationships with shoot density. Jour. Appl. Ecol. **19**: 501–517. 1982. [See also TOORN & MOOK.]
- NIERING, W. A., & R. S. WARREN. Our dynamic tidal marshes: vegetation changes as revealed by peat analysis. Conn. Arb. Bull. **12**, 22 pp. 1977.
- ONDOK, J. P. The horizontal structure of reed stands (*Phragmites communis*) and its relation to productivity. Preslia **42**: 256–261. 1970.
- PALLIS, M. The structure and history of Plav: the floating fen of the delta of the Danube. Jour. Linn. Soc. Bot. **43**: 233–290. 1916.
- PAZOUREK, J. Density of stomata in leaves of different ecotypes of *Phragmites communis*. Folia Geobot. Phytotax. Praha **8**: 15–21. 1973.
- PEARCY, R. W., J. A. BERRY, & B. BARTHOLOMEW. Field photosynthetic performance and leaf temperatures of *Phragmites communis* under summer conditions in Death Valley, California. Photosynthetica **8**: 104–108. 1974.
- RAICU, P., S. STAIU, V. STOIAN, & T. ROMAN. *Phragmites communis* Trin. chromosome complement in the Danube Delta. Hydrobiologia **39**: 83–89. 1972.
- RAKHMANINA, K. P., & Y. I. MOLOTKOVSKII. Water regime in *Phragmites australis* in southern Tadzhikistan. Sov. Jour. Ecol. **10**: 384–393. 1980.
- RAMÍREZ G., C., & N. AÑAZCO R. Variaciones estacionales en el desarrollo de *Scirpus californicus*, *Typha angustifolia*, y *Phragmites communis* en Pantanos Valdivianos, Chile. (English summary.) Agro Sur **10**: 111–123. 1982.

- SANGSTER, A. G. Electron-probe microassays for silicon in the roots of *Sorghastrum nutans* and *Phragmites communis*. *Canad. Jour. Bot.* **56**: 1074-1080. 1978.
- . Anatomical features and silica depositional patterns in the rhizomes of the grasses *Sorghastrum nutans* and *Phragmites australis*. *Ibid.* **61**: 752-761. 1983.
- SATYAMURTY, T. V. C., & V. SESHAVATHARAM. Sterility in *Phragmites communis* (Retz.) Trin. *Curr. Sci. Bangalore* **53**: 1158, 1159. 1984. [Illustration.]
- SCHIERUP, H.-H., & V. J. LARSEN. Macrophyte cycling of zinc, copper, lead, and cadmium in the littoral zone of a polluted and a non-polluted lake. I. Availability, uptake, and translocation of heavy metals in *Phragmites australis* (Cav.) Trin. *Aquatic Bot.* **11**: 197-210. 1987.
- SCHOTT, M. J., JR., & D. A. WHITE. Morphometric analysis of two adjacent populations of *Phragmites australis* in the Mississippi River Delta. *ASB Bull.* **35**: 54. 1988.
- STALTER, R. *Phragmites communis* in South Carolina. *Rhodora* **77**: 159. 1975. [First record for state.]
- STOIAN, V., A. LENOIR, P. RAICU, & R. GORENFLOT. La méiose et la taille polliniques d'individus octoploïdes du *Phragmites australis*. *Compt. Rend. Acad. Sci. Paris, D.* **278**: 457-459. 1974.
- SZAJNOWSKI, F. Relationship between leaf area index and shoot production of *Phragmites communis* Trin. *Pol. Arch. Hydrobiol.* **20**: 257-268. 1973.
- TALIGOOA, T. K., A. E. APINIS, & C. G. C. CHESTERS. Microfungi colonizing collapsed aerial parts of *Phragmites australis* Trin. in water. *Nova Hedwigia* **23**: 465-472. 1972.
- THOMPSON, D. J., & J. M. SHAY. The effects of fire on *Phragmites australis* in the Delta Marsh, Manitoba. *Canad. Jour. Bot.* **63**: 1864-1869. 1985.
- TOORN, J. VAN DER, & J. H. MOOK. The influence of environmental factors and management on stands of *Phragmites australis*. I. Effects of burning, frost, and insect damage on shoot density and shoot size. *Jour. Appl. Ecol.* **19**: 477-499. 1982.
- TORREY, J. A flora of the state of New York. 2 vols. 1056 pp. 161 pls. Albany, 1843.
- TSCHARNTKE, T. Variability of the grass *Phragmites australis* in relation to the behaviour and mortality of the gall-inducing midge *Giraudiella inclusa* (Diptera, Cecidomyiidae). *Oecologia* **76**: 504-512. 1988. [Eggs laid on the fourth internode, which is high in nutrients but low in silica.]
- VÉBER, K. Evaluation of introduced forms in experimental plantations of common reed (*Phragmites communis* Trin.). *Pol. Arch. Hydrobiol.* **22**(4): 33-42. 1980.
- WASEL, Y., & Y. RECHAW. Ecotypic differentiation in *Phragmites communis* Trin. *Hydrobiologia* **12**: 259-266. 1971.
- WALKER, J. M., & E. R. WAYGOOD. Ecology of *Phragmites communis*. I. Photosynthesis of a single shoot in situ. *Canad. Jour. Bot.* **46**: 549. 1968. [Eight percent of photosynthesis carried out by leaf sheaths.]
- WEISNER, S. E. B. Factors affecting the internal oxygen supply of *Phragmites australis* (Cav.) Trin. ex Steudel in situ. *Aquatic Bot.* **31**: 329-335. 1988. [Oxygen transport probably limits depth tolerance.]
- WEISSER, P. J., & R. J. PARSONS. Monitoring *Phragmites australis* increases from 1937 to 1976 in the Siyai Lagoon (Natal, South Africa) by means of air photo interpretation. *Bothalia* **13**: 553-556. 1981. [Area covered increased fourfold in 20 years.]
- WU, G.-L., H.-C. YE, & G.-F. LI. Embryogenic callus formation and plantlet regeneration of *Phragmites communis*. (In Chinese: English abstract.) *Acta Bot. Sinica* **29**: 361-366. 1987.
- YAMASAKI, S. Role of plant aeration in zonation of *Zizania latifolia* and *Phragmites australis*. *Aquatic Bot.* **18**: 287-297. 1984.
- & I. TANGE. Growth responses of *Zizania latifolia*, *Phragmites australis*, and *Miscanthus sacchariflorus* to varying inundation. *Aquatic Bot.* **10**: 229-239. 1981.

3. *Neyraudia* Hooker f., Fl. Brit. India 7: 305. 1896.

Caespitose perennials of damp, sunny places. Roots fibrous; rhizomes short, solid. Stems erect; nodes glabrous. Leaves several, cauline; sheath glabrous; ligule membranaceous, pilose (abaxial ligule a conspicuous cartilaginous ridge, pilose, becoming glabrous with age); blade linear-lanceolate, very slightly auriculate, the margin scabridulous, the surfaces glabrous. Inflorescences terminal (and sometimes axillary also), paniculate, much branched, plumose. Spikelets oblong, 3- to 6-flowered; glumes 2, broadly lanceolate, hyaline; rachilla internodes pilose; calluses pilose; lemmas lanceolate, 3-nerved, pilose abaxially toward the edges, the apex bifid, aristate from the notch, the awn scabridulous, stiffly excurved or recurved, sometimes slightly spiraled; paleas lanceolate, hyaline, 2-nerved. Stamens 3; anthers ellipsoid. Ovaries oblong, glabrous; styles 2, free; stigmas short, feathery. Caryopses linear, subterete, slightly dorsiventrally flattened; hilum short; embryo large. Base chromosome number 10. TYPE SPECIES: *N. madagascariensis* (Kunth) Hooker f. = *N. arundinacea* (L.) Henr. (Name an anagram of *Reynaudia*, a monotypic genus of Cuban grasses.)—SILK-REED.

Two species, both endemic to the Old World tropics. One species, *Neyraudia Reynaudiana* (Kunth) Keng, $2n = 40$, silk-reed, Burma-reed, is adventive in southern Florida (Hall; Long & Lakela; several specimens, the earliest from 1930, examined from Dade County). *Neyraudia arundinacea* (L.) Henr., $2n = 40$, a native of southern Asia, has been cultivated in southern Florida (Hall) and might be anticipated as an adventive. In *N. arundinacea* the first lemma subtends a perfect flower, while in *N. Reynaudiana* it subtends a sterile one.

Leaves of *Neyraudia* have an abaxial (external) ligule, which has been overlooked by some workers. It consists of a cartilaginous ridge convergent at its ends with the adaxial ligule. The abaxial ligule is pilose at first but usually becomes glabrous as the leaf matures.

REFERENCES:

Under subfamily references see BENTHAM (1883); CLAYTON & RENVOIZE; HALL; LONG & LAKELA; and PALMER & TUCKER.

4. *Danthonia* A. P. de Candolle in Lamarek & A. P. de Candolle, Fl. Franç. ed. 3, 3: 32. 1815, nom. cons.

Caespitose perennials. Roots fibrous or wiry. Rhizomes lacking. Stems several, unbranched, terete, more or less scabridulous, glabrous or pilose; nodes with a medial constriction. Leaves several to many, basal and cauline; sheath open, shorter than the blade, glabrous or pilose; ligule a dense fringe of short hairs; blade flat (sometimes becoming involute in age), glabrous or pilose, the margin and midvein scabridulous. Inflorescences terminal and axillary (terminal cleistogamous or chasmogamous; axillary cleistogamous, enclosed in sheath); unbranched except for 2-5 primary branches, these appressed or spreading, scabridulous. Spikelets 1-3 per branch, narrowly ellipsoid, 4- to 8-flowered; glumes 2, equal, lanceolate, (3- to) 5- to 7-nerved, acute, mucronulate, persistent; rachilla straight, glabrous; calluses pilose; lemmas elliptic,

5-nerved, pilose abaxially especially near the margin, bidentate, bearing an awn from the notch between the teeth, awn with spiraled base and straight apex (axillary florets awnless); paleas broadly elliptic. Lodicules 2 or lacking, oblong to ovate, entire or truncate. Stamens 3; anthers narrowly ellipsoid, the thecae divergent apically and basally. Ovaries ovoid; styles very short; stigmas 2, feathery. Caryopses oblong to ellipsoid, dorsiventrally flattened (the abaxial face convex, the adaxial concave), the base stipitate, the apex obtuse; embryo about $\frac{1}{3}$ as long as endosperm. Base chromosome number 12. (*Sieglingia* Bernh., nom. rejic.; including *Rytidosperma* Steudel, *Notodanthonia* Zotov.) TYPE SPECIES: *Avena spicata* L. = *D. spicata* (L.) Roemer & Schultes, typ. cons. (Named in honor of Étienne Danthoine, early nineteenth-century French botanist.)—OAT GRASS.

A genus of about 80 species (Conert, 1987), here accepted in a broad sense to include the segregate *Rytidosperma*. *Rytidosperma* was centered in Australia and New Zealand and was treated as a genus, on shaly grounds, by Clayton & Renvoize (but not by Conert, 1987). Reviewing all species attributed to *Danthonia*, Conert found no constant feature to distinguish it from *Rytidosperma*. The patterns of lemma pubescence, often used to distinguish between the two genera, show greatest diversity in Australia. Tomlinson, in an anatomical survey of the tribe, noted that *Danthonia* and *Rytidosperma* are alike in leaf-blade anatomy but differ in lodicule morphology. In *Danthonia* the lodicules lack both macrohairs and microhairs, while in *Rytidosperma* both are present. There are 32 species in Australia (Conert, 1987; Vickery) and 16 in New Zealand. There are two endemics in the Himalayas, two species in Northeast Africa, and 18 species in South America. All the North American representatives of *Danthonia* are $2n = 36$ (Gray *et al.*).

Only three species occur in the Southeast. *Danthonia spicata* (L.) Beauv. (leaf blades involute; panicle branches erect; glumes 7–14 mm long; lemma teeth (not central awn) less than 2 mm long) is widespread, occurring from Labrador to southeastern Alaska, south to western Florida and eastern Texas, and in the mountains of New Mexico and northern California. A disjunct population occurs in Hidalgo, Mexico (Conert, 1987). Canadian populations were recognized by Fernald as var. *pinetorum* Piper, said to differ in having mostly straight (not curled) basal leaves and broader, weakly nerved glumes. Dore & McNeill, however, thought most Ontario populations belonged to the typical variety.

The second southeastern species, *Danthonia compressa* Austin (leaf blades flat; panicle branches spreading; glumes 7–14 mm long; lemma teeth about 3 mm long) occurs from southern Quebec to Ohio and south to the mountains of North Carolina and Tennessee.

The third, and probably the most abundant, southeastern species is *Danthonia sericea* Nutt. (leaves involute, generally silky-pubescent; panicle branches short, erect; glumes 12–18 mm long; longer lemmas 7–10 mm long), which occurs on the Coastal Plain and Piedmont from southern New Jersey and southern Kentucky to northern Florida and Louisiana. It consists of three races, which have been accorded varietal or specific status by some workers. The

major taxonomic problem in the southeastern taxa concerns this species and its segregate *D. sericea* subsp. *epilis* (Scribner) Blomq. (*D. epilis* Scribner; leaves glabrous), reported to be restricted to the upper Piedmont of North Carolina and Georgia. Quinn & Fairbrothers made cytological preparations from 21 populations and found no karyological differences (all were $2n = 36$). Populations of *D. sericea* subsp. *epilis* in New Jersey begin growth and flowering in response to increasing soil temperature and are day-length neutral. However, populations on granitic outcrops in Georgia have their flowering time fixed genetically (Rotsettis *et al.*) and start to flower early in the year, which may be advantageous in allowing seed set before the thin soil in which the plants typically grow dries out in the summer. Gray and colleagues also studied these two regional groups of subsp. *epilis*: populations from bogs in New Jersey lacked stomata and macrohairs on the abaxial leaf surfaces but had them on the adaxial ones, while plants from well-drained sites of the Piedmont and Coastal Plain had hairs and stomata on both surfaces. Some intermediate populations from sites that were intermittently wet had abaxial hairs and stomata, but they were more plentiful on the adaxial surface. All populations had the same chromosome number, $2n = 36$. When grown in the greenhouse, both races responded to decreasing soil moisture by reduction in the width of the stomatal aperture. There is no information on compatibility or sterility between these differentiated populations, so it seems premature to recognize them taxonomically. The potential for further research is enticing.

Five other species of *Danthonia* occur in North America. *Danthonia intermedia* Vasey is boreal and cordilleran in distribution, occurring from Newfoundland to Alaska and south to the mountains of New Mexico and northern California (Cayouette & Darbyshire). It also grows on the Kamchatka Peninsula of the eastern Soviet Union (Conert, 1987; Tsvelev). *Danthonia Parryi* Scribner of the Rocky Mountains (Alberta to Colorado) is morphologically similar and probably closely related.

The second pair of western species is characterized by few-flowered inflorescences. *Danthonia californica* Bol. has an amphitropical distribution, occurring in the West from Montana to southwestern British Columbia and south to New Mexico and California, and is also reported from Chile (Munz). Pilose plants have been segregated as var. *americana* (Scribner) Hitchc. *Danthonia unispicata* (Munro ex Thurber) Munro ex Macoun has inflorescences consisting of only one (to three) spikelet(s). Its range is similar to that of *D. californica*, and Munz considered the two species as doubtfully distinct.

The Australian *Danthonia pilosa* R. Br., $2n = 48$, is adventive in California from seed planted for forage (Hitchcock, 1951; Munz).

Baum & Findlay emphasized lodicule morphology in their revision of the Canadian species of *Danthonia*. They detected four lodicule patterns, two of which are represented in the southeastern species (lodicules were absent from both *D. sericea* and *D. spicata*; club-shaped lodicules—presumably two per floret, although they did not say so—with truncate apices characterized *D. compressa*). Thus they recognized the North American species chiefly by the lodicules, a taxonomic scheme not receiving much if any subsequent acceptance. In a second paper, Findlay & Baum described a new species, *D. cana-*

densis Findlay & Baum, which occurs across Canada and the northern United States, and which differs from *D. Parryi* in the shape of the lodicules and other quantitative characters usable only by means of principal-components analysis. Such "one-character" taxonomy is problematic in a genus beset with taxonomic difficulties (see Rotsettis *et al.*; Vickery).

A notable feature of *Danthonia* is the production of cleistogamous spikelets by some species, including the three southeastern representatives. Cleistogamy was apparently first noted in the genus by Austin, who described "spikes" on short branches wholly enclosed by upper leaf sheaths. Clay (1982) investigated the reproductive biology and population genetics of *D. spicata* in North Carolina. All plants produced both axillary (cleistogamous) and aerial ("chasmogamous"—actually cleistogamous and chasmogamous) flowers,⁵ but the proportion of each kind was variable. Using vegetatively produced individuals and growing these under different conditions, Clay (1982) found that plants of the same genotype consistently produced more aerial florets in the greenhouse (40 percent) than in the field (28 percent). Larger plants produced a higher percentage of axillary florets. From genetic analysis of related individuals, he determined that 50 percent of the observed variation in production of axillary florets was genetically based and 50 percent environmentally based. Genetic differences were possibly the result of natural selection. In other natural populations the percentage of axillary florets was lower: eight or nine percent in Wyoming and zero in Michigan (Scheiner & Teeri).

Clay & Antonovics (1985b) also compared genetic variation in certain morphological characters for greenhouse- and field-grown plants of *Danthonia spicata*. They found significant genetic variation present for every character examined, both vegetative (height of flowering stem; length and width of uppermost leaf) and reproductive (number of cleistogamous flowers; length of second glume). The degree of variability among closely related individuals (plants from seeds of the same carpellate parent) suggested that the aerial florets are in part cleistogamous or self-pollinated.

The three southeastern species vary in the relative proportions of aerial and axillary florets (Clay, 1983b). *Danthonia compressa* averaged 50 percent aerial florets (minimum, 35 percent); *D. spicata* ranged from zero to 70 percent, with an average of 25 percent; and *D. sericea*, including plants assignable to *D. epilis*, averaged only five percent (maximum, ten percent). Grazing or mowing appears to favor the production of axillary florets over aerial, since such populations had the highest percentage of axillary florets.

Clay (1983b) compared the weight of diaspores (caryopsis plus clasping lemma and palea) from aerial and axillary florets in the southeastern species. In *Danthonia compressa* the average weight for both kinds was 0.80 mg, while the axillary ones were 25 percent heavier in *D. sericea* and 35 percent heavier

⁵In plants from Durham County, North Carolina, studied by Clay, all aerial florets were apparently chasmogamous. However, Clay sent seeds from his study population to Philipson, who grew them in a greenhouse in New Zealand. About half of the aerial spikelets in the resulting plants were cleistogamous (Philipson, 1986). In the discussions of genetic studies by Clay, I have used the categories aerial and axillary florets where he used "chasmogamous" and "cleistogamous." See discussion below and Philipson for further details, and see Darbyshire & Cayouette for comments.

in *D. spicata*. In the two latter species about two thirds of the difference was due to heavier lemmas and paleas in the axillary florets.

A comparison of the fitness of plants derived from aerial and axillary caryopses in *Danthonia spicata* showed only a slight difference (Clay & Antonovics, 1985a). The advantage in survival was mostly during the seed and seedling stages (Clay, 1983a). Both kinds of diaspores germinate in mid- to late spring in North Carolina. Maximum germination was about 30 percent for seeds stored dry at room temperature for several months. Axillary cleistogenes germinate over a longer period (30 percent germination in ten days) than aerial (10 to 15 percent in 30 to 40 days).

Subsequent studies by Philipson show the diversity of reproductive features in *Danthonia spicata*. She grew 49 plants (122 panicles) from a single North Carolina population and found that 52 percent of the florets from aerial panicles were in fact cleistogamous (rather than chasmogamous, as was assumed by Clay). There was wide variation in the percentage of cleistogamy, and 58 panicles (nearly 50 percent) were wholly cleistogamous. In the aerial cleistogamous florets the anthers were tiny and indehiscent; the pollen grains germinated in situ, and the tubes grew through the anther wall into the adjacent stigmas. Often, both chasmogamous and cleistogamous florets were produced in the same spikelet (the cleistogamous generally proximal to and maturing sooner than the chasmogamous). Thus, many plants produced three kinds of diaspores: chasmogamous and cleistogamous ones from aerial inflorescences, both dispersed some distance by the hygroscopic lemma awns, and atelechoric awnless axillary cleistogenes generally germinating quite close to the parent plant.

Danthonia spicata is able to grow in diverse environments, from open sandy soil in full sun to closed-canopy oak-pine forest with only ten percent of full sunlight in Michigan (Scheiner & Teeri). Individual plants have the ability to recover from severe droughts. Both genetic differentiation (microevolution or genetic drift) and phenological flexibility appear to contribute to this ecological amplitude.

Population variability based on genetics has also been demonstrated in the Australian *Danthonia caespitosa* Gaudich. (Quinn & Hodgkinson). The response to density and to temperature varied along a latitudinal gradient. The infraspecific diversity of species from different regions tends to counter the suggestion of Clayton & Renvoize that the Arundinoideae are not successful and are outside the mainstream of grass evolution.

Seed dormancy in the Australian species *Danthonia carphoides* F. Mueller ex Benth and *D. caespitosa* was investigated by Hagon. Seeds sown within two weeks of ripening had 30 percent germination. After storage at room temperature (dry) for six weeks, all dormancy was broken. Germination responses were not appreciably affected by temperatures in the 15–35°C range.

REFERENCES:

Under subfamily references see BENTHAM (1882, 1883); CLAYTON & RENVOIZE; CLEWELL; CONERT (1961; 1975; 1987); CONNOR (1987); DeWET (1954, 1956); DORE & McNEILL; HITCHCOCK (1951); MUNZ; TOMLINSON; TSVELEV; and ZOTOV.

- ABELE, K. Cytological studies in the genus *Danthonia*. Trans. Proc. Roy. Soc. S. Austral. **82**: 163–173. 1959.*
- AUSTIN, C. F. *Danthonia* DC. Bull. Torrey Bot. Club **3**: 21, 22. 1872. [Four species.]
- BAUM, B. R., & J. N. FINDLAY. Preliminary studies in the taxonomy of *Danthonia* in Canada. Canad. Jour. Bot. **51**: 437–450. 1973.
- BLAKE, S. T. *Plinthanthesis* and *Danthonia* and a review of the Australian species of *Leptochloa*. Contr. Queensland Herb. **14**: 1–19. 1972.
- BROCK, R. D., & J. A. M. BROWN. Cytotaxonomy of Australian *Danthonia*. Austral. Jour. Bot. **9**: 62–91. 1961.
- CAYOUEITE, J., & S. J. DARBYSHIRE. La répartition de *Danthonia intermedia* dans l'est du Canada. Nat. Canad. **114**: 217–220. 1988. [Distribution map; specimen citations.]
- CLAY, K. Environmental and genetic determinants of cleistogamy in a natural population of the grass *Danthonia spicata*. Evolution **36**: 734–741. 1982.
- . The differential establishment of seedlings from chasmogamous and cleistogamous flowers in natural populations of the grass *Danthonia spicata*. Oecologia **57**: 183–188. 1983a.
- . Variation in the degree of cleistogamy within and among species of the grass *Danthonia*. Am. Jour. Bot. **70**: 835–843. 1983b.
- . The effect of the fungus *Atkinsonella hypoxylon* (Clavicipitaceae) on the reproductive system and demography of the grass *Danthonia spicata*. New Phytol. **98**: 165–175. 1984.
- & J. ANTONOVICS. Demographic genetics of the grass *Danthonia spicata*: success of progeny from chasmogamous and cleistogamous flowers. Evolution **39**: 205–210. 1985a.
- & ———. Quantitative variation of progeny from chasmogamous and cleistogamous flowers in the grass *Danthonia spicata*. Ibid. 335–348. 1985b.
- & J. P. JONES. Transmission of *Atkinsonella hypoxylon* (Clavicipitaceae) by cleistogamous seed of *Danthonia spicata* (Gramineae). Canad. Jour. Bot. **62**: 2893–2895. 1984.
- & J. A. QUINN. Density of stomata and their responses to a moisture gradient in *Danthonia sericea* populations from dry and wet habitats. Bull. Torrey Bot. Club **105**: 45–49. 1978.
- CONERT, H. J. The genus *Danthonia* in Africa. Mitt. Bot. Staatssamm. München **10**: 299–308. 1971.
- . Über *Danthonia domingensis* Hackel & Pilger (Poaceae: Arundinoideae: Danthonieae). Senckenberg. Biol. **56**: 293–313. 1975.
- DARBYSHIRE, S. J. The oldest Canadian specimen in a Canadian herbarium? Pl. Press (Mississauga) **4**: 107. 1986. [*Danthonia spicata* from Newfoundland.]
- & J. CAYOUEITE. Biology of Canadian weeds. 92. *Danthonia spicata* (L.) Beauv. in Roem. & Schult. Canad. Jour. Pl. Sci. **69**: 1217–1233. 1989.
- DEWET, J. M. J. Leaf anatomy and morphology in South African species of *Danthonia*. Bothalia **7**: 303–310. 1960.
- FERNALD, M. L. Notes on *Danthonia*. Rhodora **45**: 239–246. 1943.
- FINDLAY, J. N., & B. R. BAUM. The nomenclatural implications of the taxonomy of *Danthonia* in Canada. Canad. Jour. Bot. **52**: 1573–1582. 1974. [See also BAUM & FINDLAY.]
- GRAY, J. R., J. A. QUINN, & D. E. FAIRBROTHERS. Leaf epidermis morphology in populations of the *Danthonia sericea* complex. Bull. Torrey Bot. Club **96**: 525–530. 1969.
- HAGON, M. W. Germination and dormancy of *Themeda australis*, *Danthonia* spp., *Stipa bigeniculata*, and *Bothriochloa macra*. Austral. Jour. Bot. **24**: 319–327. 1976.
- HODGKINSON, K. C., & J. A. QUINN. Adaptive variability in the growth of *Danthonia caespitosa* Gaud. populations at different temperatures. Austral. Jour. Bot. **24**: 381–396. 1976.

- & ———. Environmental and genetic control of reproduction in *Danthonia caespitosa* populations. *Ibid.* **26**: 351–364. 1978.
- PHILIPSON, M. N. A re-assessment of the form of reproduction in *Danthonia spicata* (L.) Beauv. *New Phytol.* **103**: 231–243. 1986.
- & M. C. CHRISTEY. An epiphytic/endophytic fungal associate of *Danthonia spicata* transmitted through the embryo sac. *Bot. Gaz.* **146**: 70–81. 1985. [Illustrations.]
- QUINN, J. A., & D. E. FAIRBROTHERS. Habitat, ecology and chromosome numbers of natural populations of the *Danthonia sericea* complex. *Am. Midl. Nat.* **85**: 531–536. 1971.
- & K. C. HODGKINSON. Population variability in *Danthonia caespitosa* (Gramineae) in response to increasing density under three different temperature regimes. *Am. Jour. Bot.* **70**: 1425–1431. 1983.
- ROTSETTIS, J., J. A. QUINN, & D. E. FAIRBROTHERS. Growth and flowering of *Danthonia sericea* populations. *Ecology* **53**: 227–234. 1972.
- SCHNEIDER, S. M., & C. J. GOODNIGHT. The comparison of phenotypic plasticity and genetic variation in populations of the grass *Danthonia spicata*. *Evolution* **38**: 845–855. 1984.
- , J. GUREVITCH, & J. A. TEERI. A genetic analysis of the photosynthetic properties of populations of *Danthonia spicata* that have different growth responses to light level. *Oecologia* **64**: 74–77. 1984.
- & J. A. TEERI. Phenotypic flexibility and genetic adaptation along a gradient of secondary forest succession in the grass *Danthonia spicata*. *Canad. Jour. Bot.* **64**: 739–747. 1986.
- SIMPSON, M. Value of the awn in establishing seed of *Danthonia penicillata* (Labill.) Palisot. *New Zealand Jour. Sci. Tech. A.* **1952**: 360–364. 1953.
- TOOLE, V. K. Germination of the seed of poverty grass, *Danthonia spicata*. *Jour. Am. Soc. Agron.* **31**: 954–965. 1939.*
- VICKERY, J. W. A revision of the Australian species of *Danthonia* DC. *Contr. New South Wales Natl. Herb.* **2**: 249–325. 1956.

Tribe ARISTIDEAE C. E. Hubbard in Bor, Grasses Burma Ceylon India Pakistan, 685. 1960.

5. *Aristida* Linnaeus, Sp. Pl. **1**: 82. 1753; Gen. Pl. ed. 5. 35. 1754.

Caespitose annuals or perennials of dry soils. Roots fibrous; rhizomes short or lacking. Stems several to many, erect or slightly oblique, branched from the axils, especially in the lower portion. Leaves several, cauline and basal; sheath glabrous or pilose; ligule a short, pilose membrane; external ligule (sometimes present) a cartilaginous ridge interrupted by the midvein, bearing a row of trichomes; blade linear, about as long as the sheath [absent], involute or flat, the margin and surface scabridulous. Inflorescences paniculate, more or less open. Spikelets pedicellate, 1-flowered; glumes 2, strongly unequal to subequal, sometimes awned; lemmas lanceolate, shorter or longer than the glumes, involute [convolute], 3-nerved, calluses prominent, shortly hispid, the awn conspicuous, scabridulous [plumose], with 2 lateral teeth whose bases are sometimes fused to the lower portion of the awn forming the awn column; paleas much shorter than lemmas, 2-nerved. Lodicules 2. Stamens 3 or 1; anthers linear. Ovaries shortly cylindrical; styles 2, short; stigmas plumose, laterally exserted. Caryopses compressed or terete, tightly enclosed by lemma and palea,



FIGURE 2. Spikelets or their parts. a-d, Tribe Aristideae, *Aristida longispica* (sect. *Aristida*): a, spikelet with glumes spread apart, $\times 6$; b, floret (lemma and palea) containing mature caryopsis, the 3-awned lemma completely enclosing palea and caryopsis, $\times 6$; c, caryopsis, $\times 6$; d, palea, $\times 12$. e-h, Tribe Stipeae, *Piptochaetium avenaceum* (*Stipa avenacea*): e, spikelet with single floret (only basal portion of lemma awn shown), $\times 2$; f, entire floret, showing relative length of hygroscopic lemma awn, $\times \frac{1}{2}$; g, floret (note hairy base of lemma and rachilla forming bearded, sharp-pointed callus), lemma clasping pointed palea (only base of lemma awn shown), $\times 5$; h, palea, $\times 5$. i-k, Tribe Centothecae, *Chasmanthium latifolium* (*Uniola latifolia*): i, spikelet, $\times 1\frac{1}{2}$; j, floret (lemma and palea enclosing flower), $\times 3$; k, caryopsis, $\times 5$.

sometimes longitudinally furrowed. Base chromosome number 11. TYPE SPECIES: *A. adscensionis* L., the only species included in the genus by Linnaeus. (Name from Latin *arista*, a beard or awn.)—THREE-AWN GRASS.

A genus of about 330 species distributed nearly throughout the tropical and warm-temperate regions of the world. Twenty species occur in the Southeast. The genus is reasonably well known, thanks to the worldwide studies of Henrard (1926; 1927; 1928; 1929; 1932; 1933a, b). A synopsis of the southeastern species (including those in Delaware, Maryland, Virginia, West Virginia, and Kentucky) has been presented by Allred (1986). His keys allow easy identification of the species in eastern North America.

The subgeneric classification of *Aristida* is confused and difficult. There are probably parallels in the evolution of spikelet characters noted by Henrard (1926; 1927; 1928; 1929; 1932; 1933a, b). Section *ARISTIDA* (sect. *Chaetaria* (Beauv.) Trin.) (awns of the lemma persistent) accounts for about 200 species worldwide and 18 of the 20 species in the Southeast. The section was divided by Hitchcock (1924, 1951) into informal groups, a scheme also followed by Allred (1986) and seemingly prudent in view of the lack of understanding of the phylogeny of the genus. About one third of the southeastern species are annuals (group *Dichotoma* of Hitchcock, 1951), the remainder perennials. A common situation in the section is pairs of intergrading species; for example, *A. basiramea* Engelm. ex Vasey and *A. dichotoma* Michx. Polyploidy characterizes several species of the southwestern United States (DeLisle, 1973) of the *A. purpurea* Nutt. complex (group *Purpureae* of Hitchcock, 1951). *Aristida Roemeriana* Scheele, $2n = 22$, includes only diploids; *A. longiseta* Steudel, *A. Fendleriana* Steudel, and *A. glauca* (Nees) Walp., diploids and tetraploids ($2n = 22, 44$); *A. Wrightii* Nash, diploids, tetraploids, and hexaploids; and *A. purpurea* Nutt. ($2n = 88$), octaploids as well. In none of these species were morphological criteria found to distinguish the various autopolyploids (DeLisle, 1969). The polymorphic and pantropic *A. adscensionis* L., $2n = 22$, has numerous infraspecific categories. In the New World it has been reported from Texas westward to California and southward through Central and South America.

Section *ARTHRATHERUM* (Beauv.) Reichenb. (awn column deciduous at its base) is most diverse in Africa. There are only two species in the Southeast. *Aristida desmantha* Trin. & Rupr. (longer glume less than 2 cm long; awn column 2–5(–7) mm long) occurs in our area only in Arkansas and Louisiana. *Aristida tuberculosa* Nutt. (longer glume more than 2 cm long; awn column 8–15 mm long) is widespread in the eastern United States and in all of the southeastern states except Tennessee and Arkansas. Chromosome counts (e.g., *A. brevisubulata* Maire, $2n = 22$, and *A. pallida* Steudel, $2n = 44$) suggest the occurrence of polyploidy in African representatives of this section.

The chiefly African sect. *PSEUDARTHRATHERUM* Chiov. is not represented in the New World. This section differs from the preceding in having the awn column deciduous at its summit (i.e., just below the level at which the lateral awns branch from the central one).

Also not represented in the New World is sect. *STREPTACHNE* (R. Br.) Domin

(awn column present, not articulated). This section, recognized by Lazarides and by Bourreil & Reyre, was included without comment in sect. *ARISTIDA* by Clayton & Renvoize. One of its species, *Aristida Humbertii* Bourreil & Reyre, known from Angola, is unique in the genus in that all three awns are reduced to short mucros.

REFERENCES:

- Under subfamily references see BENTHAM (1883); CLAYTON & RENVOIZE; CLEWELL; CONNOR (1987); DE WINTER; FOWLER; FOWLER & DUNLAP; HITCHCOCK (1951); and WUNDERLIN.
- ALLRED, K. W. Studies in the genus *Aristida* (Gramineae) of the southeastern United States. I. Spikelet variation in *A. purpurea*, *A. tenuispica*, and *A. virgata*. *Rhodora* **86**: 73–77. 1984a; II. Morphometric analysis of *A. intermedia* and *A. longispica*. *Ibid.* **87**: 137–145. 1985a; III. Nomenclature and a taxonomic comparison of *A. lanosa* and *A. palustris*. *Ibid.* 147–155. 1985b; IV. Key and conspectus. *Ibid.* **88**: 367–388. 1986. [Twenty species; synoptic treatment with key, distributions, illustrations of spikelets.]
- . Morphologic variation and classification of the North American *Aristida purpurea* complex (Gramineae). *Brittonia* **36**: 382–395. 1984b.
- BOGDAN, A., & A. STORRAR. Control of *Aristida* and other annuals in Kenya Rift Valley pastures. *Empire Jour. Exper. Agr.* **22**: 211–223. 1954.*
- BOURREIL, P. Réflexions sur l'écologie, la morphogénèse et l'évolution, fondées sur la culture d'*Aristida rhiniochloa*, graminée tropicale africaine. *Adansonia* II. **10**: 409–427. 1970.
- & A. GESLOT. Contribution à l'étude caryologique de diverses graminées africaines des genres *Aristida* L. et *Stipagrostis* Nees. *Adansonia* II. **11**: 125–134. 1971.
- , ———, & H. GILLET. Contribution à l'étude caryologique d'*Aristida rhiniochloa* (Graminée) d'après des spécimens d'Afrique boréale. *Adansonia* II. **11**: 685–690. 1971.
- & H. GILLET. Synthèse des connaissances et des recherches nouvelles sur *Aristida rhiniochloa*, graminée africaine amphitropicale. *Mitt. Bot. Staatssamm. München* **10**: 309–340. 1971.
- & Y. REYRE. Un nouvel aristide de l'Angola de la section *Streptachne*. *Adansonia* II. **9**: 421–427. [Includes illustrations of pollen.]
- & M. TROUIN. Contribution à l'étude caryologique de quelques aristides (Graminées) d'Afrique boréale. Conséquences taxonomiques. *Nat. Monspel. Bot.* **21**: 29–36. 1970.
- DELISLE, D. G. Chromosome number and pollen size in the genus *Aristida*. *Proc. Iowa Acad. Sci.* **86**: 74–81. 1969.
- . Chromosome numbers in the *Aristida purpurea* complex. *Southwest. Nat.* **18**: 79–83. 1973.
- HENRARD, J. T. A critical revision of the genus *Aristida*. I. *Meded. Rijks-Herb. Leiden* **54**: I–VIII, 1–220. 1926; II. *Ibid.* **54A**: 221–464. 1927; III. *Ibid.* **54B**: 465–701. 1928; Supplement. *Ibid.* **54C**: 703–747. 1933a.
- . A monograph of the genus *Aristida*. I. *Ibid.* **58**: 1–153, pls. 1–60. 1929; II. *Ibid.* **58A**: 157–325, pls. 61–169. 1932; Index. *Ibid.* **58B**: I–XII. 1933b.
- HITCHCOCK, A. S. The North American species of *Aristida*. *Contr. U. S. Natl. Herb.* **22**: 517–586. 1924.
- LAZARIDES, M. *Aristida* L. (Poaceae, Aristideae) in Australia. *Brunonia* **3**: 271–333. 1980. [Forty-seven species; discussion of problems in the genus; keys, descriptions, representative illustrations.]
- LONGHI-WAGNER, H. M. Uma nova espécie de *Aristida* L. (Gramineae) do Brasil. *Bradea* **5**: 59–62. 1988. [*A. pendula*; illustrated.]

- MATTHEI J., O. The species of *Aristida* L. (Poaceae) in Chile. *Gayana Bot.* **44**: 17–23. 1987. [Three species; illustrated.]
- SCHOLZ, H., & P. KÖNIG. Eine neue *Aristida* (Gramineae) aus Arabien. *Willdenowia* **17**: 111–113. 1988. [Illustration.]
- WARNOCK, B. H. A new three-awn grass from Trans-Pecos, Texas. *Sida* **9**: 358, 359. 1982. [*Aristida Brownii*; illustrated.]

Tribe STIPEAE Dumortier, Obs. Gram. Belg. 83. 1824.

6. *Nassella* (Trinius) Desvaux in C. Gay, *Hist. Chile Bot.* **6**: 263. 1854.

Caespitose perennials. Stems hispid or glabrous; nodes appressed-hispid, becoming glabrous with age (lower nodes usually geniculate). Leaves several to many, mostly basal; sheath pubescent or glabrous; ligule an unfringed membrane (abaxial ligule absent); blade setaceous to linear, more or less involute, hispid abaxially, less so adaxially. Inflorescences solitary, terminal, paniculate; branches flexuous, scabrid, sometimes hairy. Spikelets 1-flowered, slightly laterally compressed, disarticulation above glumes. Glumes 2, equal, broadly lanceolate, acute [awned], 3-(to 5-)nerved. Lemmas elliptic, convolute, becoming indurate, hispid basally, the surface densely covered with short, stout prickles (seemingly tuberculate), contracted into a smooth, basally fringed collar at the base of the awn, the awn 5–10 times longer than lemma, geniculate, scabrid, hairless. Paleas elliptic, much shorter than lemmas (or absent), nerveless, hyaline. Lodicules 2, glabrous, fleshy. Stamens 3; anthers oblong. Ovaries oblong, glabrous; styles free, stigmas 2. Caryopses oblong to pyriform, laterally compressed, smooth; hilum linear; embryo large, with epiblast. Base chromosome number 14. (*Urachne* sect. *Nassella* Trin. *Mém. Acad. Sci. St. Pétersb. Sci. Nat.* VI. **1**: 73. 1830.) LECTOTYPE SPECIES: *N. pungens* Desv., designated by Parodi (1947). (Origin of name unknown.)

A genus of about 50 species (Barkworth & Everett), in an area from Patagonia north through the Andes into the Caribbean region, Mexico, and the United States, and western Canada. The only southeastern representative, *Nassella leucotricha* (Trin. & Rupr.) Pohl (*Stipa leucotricha* Trin. & Rupr.), Texas wintergrass, $2n = 28$ (Gould), occurs sporadically in western Louisiana and southwestern Arkansas and ranges southward through Texas into South America. It is abundant in Texas, where it is important as a cool-season forage grass.

The caryopses of *Nassella leucotricha* have prolonged, erratic germination. They probably germinate more or less continuously throughout the year, except during lengthy dry spells (Fowler). On the basis of controlled-environment experiments, however, Call & Spoons characterized the optimal parameters for germination and hypothesized that in central Texas most germination would occur from late September through mid-November and, during mild winters, also from December through February.

REFERENCES:

Under subfamily references see BARKWORTH & EVERETT; FOWLER; FOWLER & DUNLAP; and GOULD.

BROWN, W. V. A cytological study of cleistogamous *Stipa leucotricha*. *Madroño* **10**: 97–107. 1949.

- . The relation of soil moisture to cleistogamy in *Stipa leucotricha*. Bot. Gaz. **113**: 438–444. 1952.
- CALL, C. A., & B. O. SPOONTS. Characterization and germination of chasmogamous and basal axillary cleistogamous florets of Texas wintergrass. Jour. Range Managem. **42**: 51–55. 1989.
- DYKSTERHUIS, E. J. Axillary cleistogenes in *Stipa leucotricha* and their rôle in nature. Ecology **26**: 195–199. 1945.
- PARODI, L. R. Las especies de gramíneas del género *Nassella* de la Argentina y Chile. Darwiniana **7**: 369–394. 1947. [Nine species.]

7. **Piptochaetium** J. Presl in K. Presl, Reliq. Haenk. **1**: 222. pl. 37. 1830.

Caespitose perennials of dry, open places. Stems branched only near the base, glabrous; nodes glabrous, constricted medially. Leaves several per stem; sheath ribbed, glabrous; ligule longer than wide, membranaceous (abaxial ligule absent); blade linear, flat to involute, midvein prominent, surface and margin scabridulous. Inflorescences solitary, terminal, open, paniculate; branches scabridulous. Glumes 2, equal, lanceolate-acuminate, conspicuously 3- to 5-veined, margin hyaline. Spikelets solitary, 1-flowered, terete. Calluses subulate [cuneate], covered with long, stiff, extrorse hairs. Lemmas narrowly oblong [elliptic], 3-nerved (nerves visible from adaxial surface), coriaceous, often tuberculate above, margin involute, fitting into the sulcus of the palea, apex thickened, scabridulous [spiny]; awn several times longer than lemma body, spiraled (hygroscopic), scabridulous. Paleas lanceolate, slightly longer than the lemmas, sulcate medially, glabrous, the apex protruding between the lemma margins. Lodicules 3. Stamens 3; anthers narrowly ellipsoid. Ovaries cylindrical; stigmas 2, plumose. Caryopses slenderly cylindrical, firmly enclosed by palea and lemma. Base chromosome number 11. TYPE SPECIES: *P. setifolium* J. Presl, the only species. (Name from Greek *piptein*, to fall, and *chaete*, bristle, referring to the deciduous lemma awns of the type species.)—NEEDLEGRASS.

A genus of about 30 species. The circumscription follows Barkworth & Everett, and Parodi. *Piptochaetium* is distinguished from other genera of the Stipeae by its grooved palea into which the margins of the involute lemma fit (Barkworth & Everett; Parodi). It is characterized by phloem fibers in the leaf blades, colorless cells between the abaxial epidermis and the bulliform cells, and circular to paradermally rounded vascular bundles (Parodi & Freier).

Only two species are found in the Southeast. *Piptochaetium avenaceum* (L.) Parodi (*Stipa avenacea* L.), $2n = 22, 28$, occurs from Massachusetts to southwestern Ontario (Dore & McNeill) and Michigan, south to northern Florida and eastern Texas. It is known from all the Southeastern States, where it grows in oak or pine woods, along roadsides, and in fields.

Piptochaetium avenacioides (Nash) Valencia & Costas (*Stipa avenacioides* Nash) is endemic to central Florida (Hall; Hitchcock; Wunderlin). It has longer awns (6–10 cm, vs. 4–6) and longer lemmas (12–18 mm, vs. 8–10) than *P. avenaceum* (Hall). The two species are otherwise similar and perhaps closely related, but because of considerable parallelism in the Stipeae (Barkworth & Everett), this is perhaps a premature statement. The range of *P. avenacioides*, although restricted, does not appear to overlap that of *P. avenaceum*.

REFERENCES:

Under subfamily references see BARKWORTH & EVERETT; DORE & McNEILL; HALL; HITCHCOCK (1951); PARODI; and WUNDERLIN.

PARODI, L. R., & F. FREIER. Observaciones taxonómicas sobre las gramíneas estípeas. *Ciencia Invest.* **1**: 144-146. 1945.

VALENCIA, J. I., & M. COSTAS. Estudios citotaxonómicos sobre *Piptochaetium* (Gramineae). *Bol. Soc. Argent. Bot.* **12**: 167-179. 1968.

Tribe CENTOTHECEAE Ridley, *Mater. Fl. Malay Penin.* **3**: 122. 1907.

8. *Chasmanthium* Link, *Hort. Berol.* **1**: 159. 1827.

Single-stemmed or loosely clustered perennials of open woods. Roots fibrous; rhizomes short, approximately horizontal. Stems unbranched or sparingly branched. Leaves cauline, several; sheath glabrous or pubescent; ligule a short hyaline membrane or fringe of hairs; blade linear to linear-lanceolate, scabridulous on veins, otherwise glabrous or sparsely pilose. Inflorescences terminal and/or axillary, open or contracted. Spikelets oblong, laterally flattened, sessile or pedicellate, 2- to 12-flowered; glumes 2, equal or subequal, deltoid to lanceolate, acute to acuminate, conduplicate, 3- to 7-nerved, the midvein often scabridulous; lemmas narrowly oblong, acuminate, acute, or bifid, 5- to 15-nerved, the midvein scabridulous or ciliate; paleas about as long as the lemmas, bicarinate, bulged out near the base, the keel scabridulous. Lodicules 2, lobed-truncate, 2- to 4-nerved. Stamens solitary; anther broadly oblong to linear. Ovaries oblong; styles short; stigmas 2, plumose. Caryopses ovoid to ellipsoid, laterally flattened, asymmetric, the apex subacute, the base cuneate to obtuse; embryo small, less than $\frac{1}{2}$ the length of the grain. Base chromosome number 12. (Excluding *Gouldochloa* Valdés, Morden, & Hatch.) TYPE SPECIES: *C. gracile* Link = *C. laxum* (L.) Yates, the only species included by Link. (Name from Greek *chasma*, open or gaping, and *anthos*, flower.)

A genus of five species, all endemics of eastern North America and all occurring in the Southeast. *Chasmanthium* was long included in *Uniola* (Chloridoideae). Detailed studies by Yates established the affinities of *Chasmanthium* with the Centothecae and the heterogeneity of *Uniola* as circumscribed by Hitchcock and other agrostologists. All species of *Chasmanthium* are self-compatible (Yates).

Chasmanthium latifolium (Michx.) Yates (large spikelets pedunculate, pendent), $2n = 48$, is the most distinctive species and presumably the least closely related to the others. It is found from New Jersey to Kansas south to the Florida Panhandle and south-central Texas, with outlying populations in central Nebraska and the mountains of northeastern Mexico.⁶ It occurs in all of the

⁶Reported from Manitoba by Yates as follows: Canada, Manitoba, Cow Creek Bridge, Stillwater, Gruver, 13 July 1936 (NCU). I have examined the specimen, and it is *C. latifolium* as annotated by Yates. However, it is unlikely this specimen came from Manitoba. The printed heading of the label reads "Herb. of Margaret G. Dudley/Winnipeg, Canada." There is no locality called Stillwater in Manitoba. There is a Cow Creek Bridge in Stillwater, Payne County, Oklahoma, well within the range of *C. latifolium*. Darwin Gruver is unfamiliar as a collector of Canadian plants to the Canadian botanists with whom I have corresponded. Thus I believe the specimen in question came from Oklahoma, and *C. latifolium* should be excluded from Canada's flora.

Southeastern States. Plants produce both cleistogamous and chasmogamous florets. *Chasmanthium latifolium* is tetraploid, while the other species are diploid.

The four remaining species have more or less sessile spikelets and narrow, spikelike inflorescences. The closely related *Chasmanthium sessiliflorum* (Poir-et) Yates, $2n = 24$ (southeastern Virginia to Missouri and Oklahoma, south to central Florida and eastern Texas), and *C. laxum* (L.) Yates, $2n = 24$ (southeastern New York to southeastern Virginia, Kentucky, and southeastern Missouri to central Florida and eastern Texas), are distinguished by their tiny (5–10 mm long) spikelets and purple anthers. In both species the lemma and palea spread, exposing the caryopsis at maturity; this distinguishes this species-pair from the other species of the genus, in which the caryopsis remains covered. *Chasmanthium laxum* differs from *C. sessiliflorum* only in having glabrous rather than pubescent leaf sheaths and collar.

Large sessile spikelets characterize the two remaining species, both of which have more restricted ranges than the preceding species: *Chasmanthium ornithorhynchum* (Steudel) Yates, $2n = 24$, occurs on the Gulf Coastal Plain of western Florida, southern Alabama, and southern Mississippi, to southeastern Louisiana. *Chasmanthium nitidum* (Baldwin ex Ell.) Yates, $2n = 24$, grows on the Atlantic and Gulf Coastal Plain of South Carolina, southern Georgia, and central and western Florida. In both the spikelets are 7–18 mm long, and the caryopses are covered completely by the palea and lemma. *Chasmanthium nitidum* has glabrous inflorescence axils and only one sterile lemma per spikelet, while *C. ornithorhynchum* has densely pilose axils and two to four sterile lemmas.

The hybrid *Chasmanthium laxum* \times *C. ornithorhynchum* is known from the Gulf Coast region of southern Mississippi (Yates). Hybrid populations are associated with both parental species and have enlarged sterile spikelets.

REFERENCES:

Under subfamily references see CLAYTON & RENVOIZE; CLEWELL; HALL; HITCHCOCK (1951); WUNDERLIN; and YATES.